

# Research statement

*Matthias Schütt*

## Overview

My main research interests center around algebraic and arithmetic geometry and number theory. They particularly concern questions of modularity, moduli and fields of definition. I found excellent objects of study in Calabi-Yau varieties and in elliptic surfaces. Recently I have started to extend the techniques to study other projective varieties, mainly different classes of surfaces.

This note sketches current and future research activities with references to the relevant previous achievements.

## Contents

1	Singular K3 surfaces . . . . .	1
2	Arithmetic aspects of algebraic surfaces . . . . .	2
2.1	K3 surfaces with high Picard number or extra structure . . . . .	2
2.2	Surfaces with maximal Picard number and Shimura varieties . . . . .	3
2.3	Arithmetic of Enriques surfaces . . . . .	4
3	Further projects . . . . .	4
3.1	Modularity and non-liftability of Calabi-Yau varieties . . . . .	4
3.2	Singular fibres of elliptic surfaces . . . . .	5

## 1 Singular K3 surfaces

Two-dimensional Calabi-Yau varieties are called K3 surfaces. They have not only great relevance in algebraic and arithmetic geometry, but also in string theory. My research mainly explores the deep connections between arithmetic and geometry.

A complex K3 surface  $S$  is called *singular* (in the sense of exceptional, but not non-smooth) if it has the maximal Picard number  $\rho(S) = 20$ . Hence the transcendental lattice  $T_S$  has rank two and gives rise to a two-dimensional Galois representation. If  $S$  is defined over  $\mathbb{Q}$ , modularity follows from a result by Livné [L]. My interest in singular K3 surfaces was originally motivated by the inverse problem, as raised independently by B. Mazur and D. van Straten. Recently, a joint work with N. Elkies established a positive answer:

### **Theorem 1 (Elkies, Schütt)**

*Every known newform of weight 3 with rational coefficients is associated to a singular K3 surface over  $\mathbb{Q}$ .*

Theorem 1 gives the first uniform answer to the problem of realising modular forms of weight greater than two geometrically in a single class of varieties.

As a first step towards its proof, I proved in [S4] that the associated newform is essentially determined by the CM-field  $K$  of class group exponent 1 or 2. There are 65 such fields known, and by a result of Weinberger [We], there is at most one more (which would be ruled out by the extended Riemann hypothesis). Extremal elliptic K3 surfaces and Kummer surfaces of Weil restrictions of CM-elliptic curves give singular K3 surfaces over  $\mathbb{Q}$  for 36 of these fields. Together with N. D. Elkies, I determined singular K3 surfaces over  $\mathbb{Q}$  for the remaining newforms [ES1]. To achieve this, we searched in families of elliptic K3 surfaces of high rank for suitable specialisations – a topic of independent interest that we are currently pursuing further (cf. 2.1).

My ultimate aim in this area is the classification of all singular K3 surfaces over number fields of fixed degree. In analogy to elliptic curves with CM, the number of these varieties is finite by a result of Šafarevič [Š].

### Goal

*Classify all singular K3 surface over a given number field, in particular over  $\mathbb{Q}$ , up to isomorphism.*

In [S5], I derived upper and lower bounds for the minimal degree of the field of definition. The proof uses classical results of CM and the basic theory of singular abelian and K3 surfaces, developed by Shioda with Mitani [SM] resp. Inose [SI]. For many cases, this gives already a complete solution. For the general case, I derived the following additional criterion in [S6]:

### Theorem 2

*Let  $S$  be a singular K3 surface over some number field  $L$ . Assume that  $NS(S)$  is generated by divisors defined over  $L$ . Let  $d = \text{disc}(T_S) < 0$ . Then the extension  $L(\sqrt{d})$  contains the ring class field  $H(d)$ .*

The proof of this result uses the known shape of the  $\zeta$ -function over some extension, class group theory and the Artin-Tate conjecture. Independently the theorem was proved by Elkies for the special case  $L = \mathbb{Q}$  by different techniques [E]. Theorem 2 also provides a direct proof of Šafarevič's finiteness result for singular K3 surfaces over number fields.

## 2 Arithmetic aspects of algebraic surfaces

Next to the continuation of the study of singular K3 surfaces, I have started to extend the circle of techniques and ideas to other classes of algebraic surfaces. My research currently centers on the following three themes.

### 2.1 K3 surfaces with high Picard number or extra structure

As a side-project, originating from [ES1], N. Elkies and I consider families of K3 surfaces of high Picard number. The motivating example is the Dwork pencil, consisting of one parameter-deformations of the Fermat quartic surface

$$X_\lambda : x_0^4 + x_1^4 + x_2^4 + x_3^4 = \lambda x_0 x_1 x_2 x_3.$$

We investigate this family from a purely algebraic point of view. Working with an isogenous family of K3 surface  $Y_\lambda$  (the "mirror quotient" by a  $(\mathbb{Z}/4\mathbb{Z})^2$  subgroup of

the automorphism group), we reprove that both families have Picard rank at least 19. This was originally shown by Dwork [D] using  $p$ -adic techniques. We determine the parametrising modular curve and thus find all specialisations with  $\lambda^4 \in \mathbb{Q}$  and Picard rank 20. Using Niemeier lattices, we also determine all jacobian elliptic fibrations on  $Y_\lambda$ . Similar techniques apply to other families of K3 surfaces of high Picard number.

As an application of our techniques, we want to compute the Néron-Severi lattice of (a general surface)  $X_\lambda$ . This is generally a very difficult task, but the new methods developed with T. Shioda and R. van Luijk in [SSvL] might enable us to achieve it in the present situation.

In a different direction, another project with N. D. Elkies concerns the supersingular K3 surface in characteristic two with Artin invariant one [ES2]. We classify all jacobian elliptic fibrations on this surface and relate them to each other. Such classifications have previously only been achieved in characteristic zero.

A project with R. Livné and N. Yui investigates K3 surfaces with non-symplectic automorphisms which operate trivially on the Néron-Severi group [LSY]. With minimal transcendental lattice, these surfaces have been classified by Kondō [K]. We prove their modularity and address mirror symmetry on the arithmetic side. Another paper [S7] concerns those K3 surfaces  $S$  which were not fully classified by Kondō: with an automorphism of order  $m = 2^k$  acting trivially on  $\text{NS}(S)$  while  $T_S$  has rank  $m$ . I classify these K3 surfaces completely and investigate their arithmetic. As a new feature, I compare arithmetic properties with mirror symmetry.

## 2.2 Surfaces with maximal Picard number and Shimura varieties

Surfaces with maximal Picard number and the Néron-Severi group in general have appeared in several of my papers. In characteristic zero, the Picard number  $\rho(X)$  is bounded by  $h^{1,1}(X)$ . The question arises whether the bound is attained within a fixed class of surfaces.

In the K3 case, complex surfaces with maximal Picard number are called singular (cf. Sect. 1). They lie dense in the moduli space and have interesting arithmetic structures. This is illustrated by Thm. 2 and also explains our interest in 2.1.

The general problem of maximal Picard number, however, is wide open. For instance, consider surfaces in  $\mathbb{P}^3$  of degree  $d > 4$ . So far, the only known surface with maximal Picard number was the Fermat sextic (Beauville). Recently I filled the first gap by exhibiting a quintic surface with maximal Picard number [S8]. The surface admits an automorphism of order 15 which also induces a rich arithmetic structure.

I believe more generally that automorphisms of higher order will help us to produce surfaces of higher degree with maximal Picard number, again reflecting the deep interplay between geometry and arithmetic. In order to study this, I started a project with B. van Geemen that investigates the Shimura varieties governing the variations of Hodge structures of certain algebraic surfaces. Here the concept of half twists is very powerful, since it provides a direct connection with abelian varieties by [vG]. The abelian varieties can be used to study the degeneration locus on the Shimura variety.

These methods can also be applied to K3 surfaces. Namely, we plan to extend the study of the families of K3 surfaces in [S7] from the point of view of Hodge structures and Shimura curves.

## 2.3 Arithmetic of Enriques surfaces

With K. Hulek, I am currently developing a program for arithmetic properties of Enriques surfaces. The main problems concern the question of fields of definition and the Galois action on the Néron-Severi group. There is great evidence that the rich theory of K3 surface will give rise to interesting phenomena on the side of Enriques surfaces.

As starting objects, we are particularly considering Kummer surfaces and special Enriques surfaces, such as those with finite automorphism groups and those which are covered by singular K3 surfaces.

An important line of argument involves elliptic fibrations. Namely, we can start with a rational elliptic surface and consider quadratic base changes (which generally are K3 surfaces). We found a canonical way to endow the base change with Enriques involutions. Geometrically this makes a recent result of Ohashi explicit that K3 surfaces may admit arbitrarily many Enriques involutions [O].

Specifically, this technique applies to singular K3 surfaces where we can construct Enriques involutions explicitly in an algebraic way (without using lattice theory). In fact, we find that a singular K3 surface admits an Enriques involution if and only if it admits an Enriques involution of the above type. In consequence, arithmetic results about singular K3 surfaces as in section 1 can be descended to the Enriques quotients. Overall we obtain a picture that is very similar to section 1:

### **Theorem 3 (Hulek, Schütt)**

*Let  $Y$  denote a complex Enriques surface that is covered by a singular K3 surface  $S$ . Let  $d$  denote the discriminant of  $NS(S)$ . Then  $Y$  admits a model over the ring class field  $H(d)$ .*

We aim to investigate this uniform result further. Possible extensions direct towards Thm. 2 and the classification of Enriques surfaces as above over fixed number fields.

## 3 Further projects

### 3.1 Modularity and non-liftability of Calabi-Yau varieties

In [S1] und [S2] I found several modular Calabi-Yau threefolds over  $\mathbb{Q}$ . Following a construction of Schoen [Sc], these varieties arose as fibre products of semi-stable rational elliptic surfaces with section. Recently Cynk and van Straten showed how modularity gives rise to non-liftable threefolds at the bad primes [CvS]. This cannot happen for Calabi-Yau varieties in dimension one and two by a result of Deligne.

In a new project with S. Cynk, we are going to extend these ideas to other fibre products [CS]. We allow the rational elliptic surfaces to involve additive singular fibres in such a way that the fibre product still admits a small resolution in the category of algebraic spaces. The results will be innovative in two respects: on the one hand introducing new modular varieties, on the other hand showing non-liftability at new primes. As an illustration, we mention the following implications of our work:

**Theorem 4**

- (i) *There is a non-liftable three-dimensional Calabi-Yau space in any characteristic less than 100 except possibly for 53, 83.*
- (ii) *There is a non-liftable three-dimensional Calabi-Yau space in characteristic 521201.*

**3.2 Singular fibres of elliptic surfaces**

The singular fibres of elliptic surfaces have been classified by Kodaira and Tate. The question arises what the maximal singular fibres within a fixed class of elliptic surfaces are. Here it is convenient to consider elliptic surfaces over  $\mathbb{P}^1$  with section and fix the Euler number. Naturally, the situation differs in characteristic zero and  $p > 0$ : The latter features supersingular surfaces with  $\rho = b_2$  while over  $\mathbb{C}$ , Lefschetz' theorem gives  $\rho \leq h^{1,1}$ . The papers [S3] and [SS1] (together with A. Schweizer) derive a complete solution when the maximal fibres mod  $p$  exceed the restriction from Lefschetz' bound in characteristic zero. In [SS2], we also proved uniqueness in the K3 case.

We want to continue these investigations in a new project concerning elliptic surfaces over an elliptic base curve with only one or two singular fibres. Over the ground field  $\mathbb{C}$ , these have been completely classified. In positive characteristic, we aim to establish essentially the same classification; only small characteristics come with some extra features due to bad reduction and wild ramification.

**References**

- [CS] Cynk, S., Schütt, M.: *Non-liftable Calabi-Yau spaces*, in preparation.
- [CvS] Cynk, S., van Straten, D.: *Small Resolutions and Non-Liftable Calabi-Yau three-folds*, manus. math. **130** (2009), 233–249.
- [D] Dwork, B.: *p-adic cycles*, Publ. math. I.H.E.S. **37** (1969), 27–115.
- [E] Elkies, N. D.: *The maximal Mordell-Weil rank of an elliptic K3 surface over  $\mathbb{Q}(t)$* , talk at conference on *Birational Automorphisms of Compact Complex Manifold and Dynamical Systems* at Nagoya University, Aug 28, 2007.
- [ES1] Elkies, N. D., Schütt, M.: *Modular forms and K3 surfaces*, preprint (2008), arXiv: 0809.0830.
- [ES2] Elkies, N. D., Schütt, M.: *Jacobian elliptic fibrations on the supersingular K3 surface in characteristic two with Artin invariant one*, in preparation.
- [vG] van Geemen, B.: *Half twists of Hodge structures of CM-type*, J. Math. Soc. Japan **53** (2001), no. 4, 813–833.
- [K] Kondō, S.: *Automorphisms of algebraic K3 surfaces which act trivially on Picard groups*, J. Math. Soc. Japan **44**, No. 1 (1992), 75–98.
- [L] Livné, R.: *Motivic Orthogonal Two-dimensional Representations of  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$* , Israel J. of Math. **92** (1995), 149–156.
- [LSY] Livné, R., Schütt, M., Yui, N.: *The modularity of K3 surfaces with non-symplectic group actions*, preprint (2009), arXiv: 0904.1922.
- [O] Ohashi, H.: *On the number of Enriques quotients of a K3 surface*, Publ. RIMS **43** (2007), no. 1, 181–200.
- [Š] Šafarevič, I. R.: *On the arithmetic of singular K3-surfaces*, in: *Algebra and analysis* (Kazan 1994), de Gruyter (1996), 103–108.
- [Sc] Schoen, C.: *On Fiber Products of Rational Elliptic Surfaces with Section*, Math. Z. **197** (1988), 177–199.

- [S1] Schütt, M.: *New examples of modular rigid Calabi-Yau threefolds*, Collect. Math. **55**, 2 (2004), 219–228.
- [S2] Schütt, M.: *On the modularity of three Calabi-Yau threefolds with bad reduction at 11*, Canad. Math. Bull. **49** (2006), 296–312.
- [S3] Schütt, M.: *The maximal singular fibres of elliptic K3 surfaces*, Arch. Math. **87**, Nr. 4 (2006), 309–319.
- [S4] Schütt, M.: *CM newforms with rational coefficients*, Ramanujan J. **19** (2009), 187–205.
- [S5] Schütt, M.: *Fields of definition of singular K3 surfaces*, Communications in Number Theory and Physics **1**, 2 (2007), 307–321.
- [S6] Schütt, M.: *K3 surfaces of Picard rank 20 over  $\mathbb{Q}$* , preprint (2008), arXiv: 0804.1558.
- [S7] Schütt, M.: *K3 surfaces with non-symplectic automorphism of 2-power order*, to appear in Journal of Algebra, preprint (2008), arXiv: 0807.3708.
- [S8] Schütt, M.: *A quintic surface with maximal Picard number*, preprint (2008), arXiv: 0812.3519.
- [SS1] Schütt, M., Schweizer, A.: *Davenport-Stothers inequalities and elliptic surfaces in positive characteristic*, Quarterly J. Math. **59** (2008), 499–522.
- [SS2] Schütt, M., Schweizer, A.: *On the uniqueness of elliptic K3 surfaces with maximal singular fibre*, preprint (2007), arXiv: 0712.3873.
- [SSvL] Schütt, M., Shioda, T., van Luijk, R.: *Lines on Fermat surfaces*, preprint (2008), arXiv: 0812.2377.
- [SI] Shioda, T., Inose, H.: *On Singular K3 Surfaces*, in: W. L. Baily Jr., T. Shioda (eds.), *Complex analysis and algebraic geometry*, Iwanami Shoten, Tokyo (1977), 119–136.
- [SM] Shioda, T., Mitani, N.: *Singular abelian surfaces and binary quadratic forms*, in: *Classification of algebraic varieties and compact complex manifolds*, SLN **412** (1974), 259–287.
- [We] Weinberger, P. J.: *Exponents of the class groups of complex quadratic fields*, Acta Arith. **22** (1973), 117–124.

Matthias Schütt,  
Institut für Algebraische Geometrie  
Leibniz Universität Hannover  
Welfengarten 1  
30167 Hannover  
Germany  
schuett@math.uni-hannover.de