

(Math 170) Homework 4:

Due February 16, 2007

Exercise 1: Find the following up to 6 decimal places

- (a) $[1]$, $[1, 4]$, $[1, 4, 1]$ and $[1, 4, 1, 4]$.
- (b) $[1]$, $[1, 3]$, $[1, 3, 1]$ and $[1, 3, 1, 3]$.
- (c) Which is larger?

Exercise 2: Find an expression of the form a/b where a, b are real numbers for

- (a) $[1, 4, 1, 4, \dots]$
- (b) $[1, 3, 1, 3, \dots]$

Let

$$q_n = \sum_{k=1}^n \frac{1}{k!}$$

Exercise 3: Prove that $\{q_n\}_{n \in \mathbb{N}}$ is a Cauchy Sequence.

Exercise 4: Let $r = \lim_{n \rightarrow \infty} q_n$. Approximate r to 6 decimal places.

Let $\{q_n\}_{n \in \mathbb{N}}$, $\{p_n\}_{n \in \mathbb{N}}$ be Cauchy Sequences which are equivalent. Further let $\{a_n\}_{n \in \mathbb{N}}$, $\{b_n\}_{n \in \mathbb{N}}$ also be Cauchy Sequences which are equivalent.

Exercise 5: (a) Show, using properties of sums of Cauchy Sequences, that $\{q_n - a_n\}_{n \in \mathbb{N}}$ is equivalent to $\{p_n - b_n\}_{n \in \mathbb{N}}$.
 (a) Show, without using properties of sums of Cauchy sequences discussed in class, that $\{q_n - a_n\}_{n \in \mathbb{N}}$ is equivalent to $\{p_n - b_n\}_{n \in \mathbb{N}}$.

Exercise 6: (2 Bonus Points) Show $\{q_n \cdot a_n\}_{n \in \mathbb{N}}$ is equivalent to $\{p_n \cdot b_n\}_{n \in \mathbb{N}}$