

CANONICAL FORKING IN AECS

WILL BONEY, RAMI GROSSBERG, ALEXEI KOLESNIKOV, AND SEBASTIEN VASEY

ABSTRACT. Boney and Grossberg [BG] proved that every nice AEC has an independence relation. We prove that this relation is unique: in any given AEC, there can exist at most one independence relation that satisfies existence, extension, uniqueness and local character. While doing this, we study more generally the properties of independence relations for AECs and also prove a canonicity result for Shelah's good frames. The usual tools of first-order logic (like the finite equivalence relation theorem or the type amalgamation theorem in simple theories) are not available in this context. In addition to the loss of the compactness theorem, we have the added difficulty of not being able to assume that types are sets of formulas. We work axiomatically and develop new tools to understand this general framework.

CONTENTS

1. Introduction	2
2. Notation and prerequisites	3
3. Independence relations	6
4. Comparing two independence relations	13
5. Relationship between various properties	18
6. Applications	25
7. Conclusion	30
References	31

Date: March 4, 2016

AMS 2010 Subject Classification: Primary: 03C48, 03C45 and 03C52. Secondary: 03C55, 03C75, 03C85 and 03E55.

Key words and phrases. Abstract Elementary Classes; forking; Classification Theory; stability; good frames.

This material is based upon work done while the first author was supported by the National Science Foundation under Grant No. DMS-1402191.

This material is based upon work done while the fourth author was supported by the Swiss National Science Foundation under Grant No. 155136.

1. INTRODUCTION

Let K be an abstract elementary class (AEC) which satisfies amalgamation, joint embedding, and which does not have maximal models. These assumptions allow us to work inside its monster model \mathfrak{C} . The main results of this paper are:

- (1) There is at most one independence relation satisfying existence, extension, uniqueness and local character (Corollary 5.19).
- (2) Under some reasonable conditions, the coheir relation of [BG] has local character and is canonical (Theorems 6.4 and 6.7).
- (3) Shelah’s weakly successful good λ -frames are canonical: an AEC can have at most one such frame (Theorem 6.13).

To understand the relevance of the results, some history is necessary.

In 1970, Shelah discovered the notion “ $\text{tp}(\bar{a}/B)$ forks over A ” (for $A \subseteq B$), a generalization of Morley’s rank in ω -stable theories. Its basic properties were published in [She78].

In 1974, Lascar [Las76, Theorem 4.9] established that for superstable theories, any relation between \bar{a} , B , A satisfying the basic properties of forking is Shelah’s forking relation. In 1984, Harnik and Harrington [HH84, Theorem 5.8] extended Lascar’s abstract characterization to stable theories. Their main device was the finite equivalence relation theorem. In 1997, Kim and Pillay [KP97, Theorem 4.2] published an extension to simple theories, using the independence theorem (also known as the type-amalgamation theorem).

This paper deals with the characterization of independence relations in various non-elementary classes. An early attempt on this problem can be found in Kolesnikov’s [Kol05], which focuses on some important particular cases (e.g. homogeneous model theory and classes of atomic models). We work in a more general context, and only rely on the abstract properties of independence. We cannot assume that types are sets of formulas, so work only with Galois (i.e. orbital) types.

In [She87, Chapter II] (which later appeared as [She09b, Chapter V.B]), Shelah gave the first axiomatic definition of independence in AECs, and showed that it generalized first-order forking. In [She09a, Chapter II], Shelah gave a similar definition, localized to models of a particular size λ (the so-called “good λ -frame”). Shelah proved that a good frame existed, under very strong assumptions (typically, the class is required to be categorical in two consecutive cardinals).

Recently, working with a different set of assumptions (the existence of a monster model and tameness), Boney and Grossberg [BG] gave conditions (namely a form of Galois stability and the extension property for coheir) under which an AEC has a global independence relation. This showed that one could study independence in

a broad family of AECs. Our paper is strongly motivated by both [She09a, Chapter II] and [BG].

The paper is structured as follows. In Section 2, we fix our notation, and review some of the basic concepts in the theory of AECs. In Section 3, we introduce independence relations, the main object of study of this paper, as well as some important properties they could satisfy, such as extension and uniqueness. We consider two examples: coheir and nonsplitting.

In Section 4, we prove a weaker version of (1) (Corollary 4.14) that has some extra assumptions. This is the core of the paper.

In Section 5, we go back to the properties listed in Section 3 and investigate relations between them. We show that some of the hypotheses in Corollary 4.14 are redundant. For example, we show that the symmetry and transitivity properties follow from existence, extension, uniqueness, and local character. We conclude by proving (1). Finally, in Section 6, we apply our methods to the coheir relation considered in [BG] and to Shelah's good frames, proving (2) and (3).

While we work in a more general framework, the basic results of Sections 2-3 often have proofs that are very similar to their first-order analogs. Readers feeling confident in their knowledge of first-order nonforking can start reading directly from Section 4 and refer back to Sections 2-3 as needed.

This paper was written while the first and fourth authors were working on a Ph.D. under the direction of Rami Grossberg at Carnegie Mellon University. They would like to thank Professor Grossberg for his guidance and assistance in their research in general and in this work specifically.

An early version of this paper was circulated already in early 2014. Since that time, Theorem 5.14 has been used by the fourth author to build a good frame from amalgamation, tameness, and categoricity in a suitable cardinal [Vasa]. With VanDieren, the fourth author has also used it to deduce a certain symmetry property for nonsplitting in classes with amalgamation categorical in a high-enough cardinal [VV], with consequences on the uniqueness of limit models. The question of canonicity of forking in more local setups (e.g. when the independence relation is only defined for certain types over models of a certain size) is pursued further in [Vasb]. The latter preprint addresses Questions 5.5, 6.14, 7.1, and 7.2 posed in this paper.

2. NOTATION AND PREREQUISITES

We assume the reader is familiar with abstract elementary classes and the basic related concepts. We briefly review what we need in this paper, and set up some notation.

Hypothesis 2.1. We work in a fixed abstract elementary class (K, \prec) which satisfies amalgamation and joint embedding, and has no maximal models.

2.1. The monster model.

Definition 2.2. Let $\mu > \text{LS}(K)$ be a cardinal. For models $M \prec N$, we say N is a μ -universal extension of M if for any $M' \succ M$, with $\|M'\| < \mu$, M' can be embedded inside N over M , i.e. there exists a K -embedding $f : M' \rightarrow N$ fixing M pointwise. We say N is a universal extension of M if it is a $\|M\|^+$ -universal extension of M .

Definition 2.3. Let $\mu > \text{LS}(K)$ be a cardinal. We say a model N is μ -model homogeneous if for any $M \prec N$, N is a μ -universal extension of M . We say M is μ -saturated if it is μ -model homogeneous (this is equivalent to the classical definition by [She01, Lemma 0.26]).

Definition 2.4 (Monster model). Since K has amalgamation and joint embedding properties and has no maximal models, we can build a strictly increasing continuous chain $(\mathfrak{C}_i)_{i \in \text{OR}}$, where for all i , \mathfrak{C}_{i+1} is universal over \mathfrak{C}_i . We call the union $\mathfrak{C} := \bigcup_{i \in \text{OR}} \mathfrak{C}_i$ the monster model¹ of K .

Any model of K can be embedded inside the monster model, so we will adopt the convention that *any set or model we consider is a subset or a substructure of \mathfrak{C}* .

We write $\text{Aut}_A(\mathfrak{C})$ for the set of automorphisms of \mathfrak{C} fixing A pointwise. When $A = \emptyset$, we omit it.

We will use the following without comments.

Remark 2.5. Let M, N be models. By our convention, $M \prec \mathfrak{C}$ and $N \prec \mathfrak{C}$, thus by the coherence axiom, $M \subseteq N$ implies $M \prec N$.

Definition 2.6. Let I be an index set. Let $\bar{A} := (A_i)_{i \in I}$, $\bar{B} := (B_i)_{i \in I}$ be sequences of sets, and let C be a set. We write $f : \bar{A} \equiv_C \bar{B}$ to mean that $f \in \text{Aut}_C(\mathfrak{C})$, and for all $i \in I$, $f[A_i] = B_i$. We write $\bar{A} \equiv_C \bar{B}$ to mean that $f : \bar{A} \equiv_C \bar{B}$ for some f . When C is empty, we omit it.

We will most often use this notation when I has a single element, or when all the sets are singletons. In the later case, we identify a set with the corresponding singleton, i.e. if $\bar{a} = (a_i)_{i \in I}$ and $\bar{b} := (b_i)_{i \in I}$ are sequences, we write $f : \bar{a} \equiv_C \bar{b}$ instead of $f : \bar{A} \equiv_C \bar{B}$, with $A_i := \{a_i\}$, $B_i := \{b_i\}$. We write $\text{gtp}(\bar{a}/C)$ for the \equiv_C equivalence class of \bar{a} . This corresponds to the usual notion of Galois types first defined in [She01, Definition 0.17].

Note that for sets A, B , we have $f : A \equiv_C B$ precisely when there are enumerations \bar{a}, \bar{b} of A and B respectively such that $f : \bar{a} \equiv_C \bar{b}$.

¹Since \mathfrak{C} is a proper class, it is strictly speaking not an element of K . We ignore this detail, since we could always replace OR in the definition of \mathfrak{C} by a cardinal much bigger than the size of the models under discussion.

2.2. Tameness and stability. Although we will make no serious use of it in this paper, we briefly review the notion of tameness. While it appears implicitly in [She99], tameness was first introduced in [GV06b] and used in [GV06a] to prove an upward categoricity transfer. Our definition follows [Bon14b, Definition 3.1].

Definition 2.7 (Tameness). Let $\kappa > \text{LS}(K)$. Let α be a cardinal. We say K is κ -tame for α -length types if for any tuples \bar{a}, \bar{b} of length α , and any $M \in K$, if $\bar{a} \not\equiv_M \bar{b}$, there exists $M_0 \prec M$ of size $\leq \kappa$ such that $\bar{a} \not\equiv_{M_0} \bar{b}$.

We say K is $(< \kappa)$ -tame for α -length types if for any tuples \bar{a}, \bar{b} of length α , and any $M \in K$, if $\bar{a} \not\equiv_M \bar{b}$, there exists $M_0 \prec M$ of size $< \kappa$ such that $\bar{a} \not\equiv_{M_0} \bar{b}$.

We say K is κ -tame if it is κ -tame for 1-length types. We say K is *fully* κ -tame if it is κ -tame for all lengths. Similarly for $(< \kappa)$ -tame.

The following dual of tameness is introduced in [Bon14b, Definition 3.3]:

Definition 2.8 (Type shortness). Let $\kappa > \text{LS}(K)$. Let μ be a cardinal. We say K is κ -type short over μ -sized models if for any index set I , any enumerations $\bar{a} := (a_i)_{i \in I}$, $\bar{b} := (b_i)_{i \in I}$ of type I , and any $M \in K_\mu$, if $\bar{a} \not\equiv_M \bar{b}$, there is $I_0 \subseteq I$ of size $\leq \kappa$ such that $\bar{a}_{I_0} \not\equiv_M \bar{b}_{I_0}$. Here $\bar{a}_{I_0} := (a_i)_{i \in I_0}$.

We define $(< \kappa)$ -type short over μ -sized models similarly.

We say K is *fully* κ -type short if it is κ -type short over μ -sized models for all μ . Similarly for $(< \kappa)$ -type short.

We also recall that we can define a notion of stability:

Definition 2.9 (Stability). Let $\lambda \geq \text{LS}(K)$ and α be cardinals. We say K is α -stable in λ if for any $M \in K_\lambda$, $S^\alpha(M) := \{\text{gtp}(\bar{b}/M) \mid \bar{b} \in {}^\alpha \mathfrak{C}\}$ has cardinality $\leq \lambda$. Equivalently, given any collection $\{A_i\}_{i < \lambda^+}$, where for all $i < \lambda^+$, $|A_i| = \alpha$, there exists $i < j$ such that $A_i \equiv_M A_j$.

We say K is *stable* in λ if it is 1-stable in λ .

We say K is α -stable if it is α -stable in λ for some $\lambda \geq \text{LS}(K)$. We say K is *stable* if it is 1-stable in λ for some $\lambda \geq \text{LS}(K)$. We write “unstable” instead of “not stable”.

Remark 2.10. If $\alpha < \beta$, and K is β -stable in λ , then K is α -stable in λ .

The following follows from [Bon, Theorem 3.1].

Fact 2.11. Let $\lambda \geq \text{LS}(K)$. Let α be a cardinal. Assume K is stable in λ and $\lambda^\alpha = \lambda$. Then K is α -stable in λ .

3. INDEPENDENCE RELATIONS

In this section, we define independence relations, the main object of study of this paper. We then consider two examples: coheir and nonsplitting.

3.1. Basic definitions.

Definition 3.1 (Independence relation). An *independence relation* \perp is a set of triples of the form (A, M, N) where A is a set, M, N are models (i.e. $M, N \in K$), $M \prec N$. Write $A \perp_M N$ for $(A, M, N) \in \perp$. When $A = \{a\}$, we may write $a \perp_M N$ for $A \perp_M N$. We require that \perp satisfies the following properties:

- (I) Invariance: Assume $(A, M, N) \equiv (A', M', N')$. Then $A \perp_M N$ if and only if $A' \perp_{M'} N'$.
- (M) Left and right monotonicity: If $A \perp_M N$, $A' \subseteq A$, $M \prec N' \prec N$, then $A' \perp_M N'$.
- (B) Base monotonicity: If $A \perp_M N$, and $M \prec M' \prec N$, then $A \perp_{M'} N$.

We write \perp_M for \perp restricted to the base set M , and similarly for e.g. $A \perp_M$.

In what follows, \perp always denotes an independence relation.

Remark 3.2. To avoid relying on a monster model, we could introduce an ambient model \widehat{N} as a fourth parameter in the above definition (i.e. we would write $A \perp_M^{\widehat{N}} N$). This would match the approach in [She09b, Chapter V.B] and [She09a, Chapter II] where the existence of a monster model is not assumed. We would require that \widehat{N} contains the other parameters A, M and N . To avoid cluttering the notation, we will not adopt this approach, but generalizing most of our results to this context should cause no major difficulty. Some simple cases will be treated in the discussion of good frames in Section 6. In [Vasb], many of the results of this paper are stated in a “monsterless” framework.

We will consider the following properties of independence²:

- $(C)_\kappa$ Continuity: If $A \not\perp_M N$, then there exists $A^- \subseteq A$, $B^- \subseteq N$ of size strictly less than κ such that for all $N_0 \succ M$ containing B^- , $A^- \not\perp_M N_0$.

²Continuity, transitivity, uniqueness, existence and extension are adapted from [MS90]. Symmetry comes from [She09a, Chapter II].

- (T) Left transitivity: If $M_1 \downarrow_{M_0} N$, and $M_2 \downarrow_{M_1} N$, with $M_0 \prec M_1 \prec M_2$, then $M_2 \downarrow_{M_0} N$.
- (T_{*}) Right transitivity: If $A \downarrow_{M_0} M_1$, and $A \downarrow_{M_1} M_2$, with $M_0 \prec M_1 \prec M_2$, then $A \downarrow_{M_0} M_2$.
- (S) Symmetry: If $A \downarrow_M N$, then there is $M' \succ M$ with $A \subseteq M'$ such that $N \downarrow_{M'} M'$. If A is a model extending M , one can take $M' = A$ ³.
- (U) Uniqueness: If $A \downarrow_M N$, $A' \downarrow_M N$, and $f : A \equiv_M A'$, then $g : A \equiv_N A'$ for some g so that $g \upharpoonright A = f \upharpoonright A$.
- (E) The following properties hold:
 - (E₀) Existence: for all sets A and models M , $A \downarrow_M M$.
 - (E₁) Extension: Given a set A , and $M \prec N \prec N'$, if $A \downarrow_M N$, then there is $A' \equiv_N A$ such that $A' \downarrow_{M'} N'$.
- (L) Local character: $\kappa_\alpha(\downarrow) < \infty$ for all α , where $\kappa_\alpha(\downarrow) := \min\{\lambda \in \text{REG} \cup \{\infty\} : \text{for all } \mu = \text{cf } \mu \geq \lambda, \text{ all increasing, continuous chains } \langle M_i : i \leq \mu \rangle \text{ and all sets } A \text{ of size } \alpha, \text{ there is some } i_0 < \mu \text{ so } A \downarrow_{M_{i_0}} M_\mu\}$.
- (E₊) Strong extension: A technical property used in the proof of canonicity. See Definition 4.4.

For (P) a property that is not local character, and M a model, when we say \downarrow has (P)_M, we mean \downarrow has (P) (i.e. \downarrow has (P) when the base is restricted to be M). If P is either (T) or (T_{*}), (P)_M means we assume $M_0 = M$ in the definition.

Whenever we are considering two independence relations $\downarrow^{(1)}$ and $\downarrow^{(2)}$, we write (P⁽¹⁾) as a shorthand for “ $\downarrow^{(1)}$ has (P)”, and similarly for (P⁽²⁾).

Notice the following important consequence of (E):

Remark 3.3. Assume \downarrow has (E)_M. Then for any A , and $N \succ M$, there is $A' \equiv_M A$ such that $A' \downarrow_M N$ (use (E₀)_M to see $A \downarrow_M M$, and then use (E₁)_M).

Assuming (T_{*})_M, this last statement is actually equivalent to (E)_M.

The property (E₊) will be introduced and motivated later in the paper. For now, we note that there is an asymmetry in our definition of an independence relation: the parameter on the left is allowed to be an arbitrary set, while the parameter on the right must be a model extending the base. This is because we have in mind

³This second part actually follows from monotonicity and the first part.

the analogy “ $a \downarrow_M N$ if and only if $\text{tp}(a/N)$ does not fork over M ”, and in AECs, types over models are much better behaved than types over sets.

The price to pay is that the statement of symmetry is not easy to work with. Assume for example we know an independence relation satisfies (T) and (S) . Should it satisfy (T_*) ? Surprisingly, this is not easy to show. We prove it in Lemma 5.9, assuming (E) . For now, we prepare the ground by showing how to extend an independence relation to take arbitrary sets on the right hand side.

Definition 3.4 (Closure of an independence relation). We call $\bar{\downarrow}$ a *closure* of \downarrow if $\bar{\downarrow}$ is a relation defined on all triples of the form (A, M, B) , where M is a model (but maybe $M \not\subseteq B$). We require it satisfies the following properties:

- For all A , and all $M \prec N$, $A \downarrow_M N$ if and only if $A \bar{\downarrow}_M N$.
- (I) Invariance: If $(A, M, B) \equiv (A', M', B')$, then $A \bar{\downarrow}_M B$ if and only if $A' \bar{\downarrow}_{M'} B'$.
- (M) Left and right monotonicity: If $A \bar{\downarrow}_M B$ and $A' \subseteq A$, $B' \subseteq B$, then $A' \bar{\downarrow}_M B'$.
- (B) Base monotonicity: If $A \bar{\downarrow}_M B$, and $M \prec M' \subseteq M \cup B$, then $A \bar{\downarrow}_{M'} B$.

The *minimal closure* of \downarrow is the relation $\bar{\downarrow}$ defined by $A \bar{\downarrow}_M C$ if and only if there exists $N \succ M$, with $C \subseteq N$, so that $A \downarrow_M N$.

It is straightforward to check that the minimal closure of \downarrow is the smallest closure of \downarrow but there might be others (and they also sometimes turn out to be useful), see the coheir and explicit nonsplitting examples below.

We can adapt the list of properties to a closure $\bar{\downarrow}$.

Definition 3.5.

- We say $\bar{\downarrow}$ has (S) if for all sets A, B , $A \bar{\downarrow}_M B$ if and only if $B \bar{\downarrow}_M A$.
- We say that $\bar{\downarrow}$ has $(C)_\kappa$ if whenever $A \bar{\downarrow}_M B$, there exists $A^- \subseteq A$, $B^- \subseteq B$ of size strictly less than κ such that $A^- \not\bar{\downarrow}_M B^-$.

- We say that $\bar{\downarrow}$ has (E_1) if whenever $A \bar{\downarrow}_M C$, and $C \subseteq C'$, there exists $A' \equiv_{MC} A$ such that $A' \bar{\downarrow}_M C'$.
- We say that $\bar{\downarrow}$ has (U) if whenever $A \bar{\downarrow}_M C$, $A' \bar{\downarrow}_M C$, and $f : A \equiv_M A'$, there is $g : A \equiv_{MC} A'$ with $g \upharpoonright A = f \upharpoonright A$.
- We say that $\bar{\downarrow}$ has (T) if whenever $M_0 \prec M_1 \prec M_2$, $M_2 \bar{\downarrow}_{M_1} C$, and $M_1 \bar{\downarrow}_{M_0} C$, we have $M_2 \bar{\downarrow}_{M_0} C$.
- The statements of (T_*) , (E_0) , (L) are unchanged. We will not need to use (E_+) on a closure.

For an arbitrary closure, we cannot say much about the relationship between the properties satisfied by \downarrow and those satisfied by $\bar{\downarrow}$. The situation is different for the minimal closure, but we defer our analysis to section 5.

Remark 3.6. Shelah’s notion of a good λ -frame introduced in [She09a, Chapter II] is another axiomatic approach to independence in AECs. There are several key differences with our framework. In particular, good λ -frames only operate on λ -sized models and singleton sets. On the other hand, the theory of good λ -frames is very developed; see e.g. [She09a, JS12, JS13].

An earlier framework which is closer to our own is the “Existential framework” AxFr_3 (see [She09b, Definition V.B.1.9]). The key differences are that AxFr_3 only defines $M_1 \bar{\downarrow}_M^{\hat{N}} M_2$ when $M \prec M_\ell$, $\ell = 1, 2$, AxFr_3 (essentially) assumes $(C)_{\aleph_0}$, while we seldom need continuity, and local character (a property crucial to our canonicity proof) is absent from the axioms of AxFr_3 .

3.2. Examples. Though so far developed abstractly, this framework includes many previously studied independence relations.

Definition 3.7 (Coheir, [BG]). Fix a cardinal $\kappa > \text{LS}(K)$. We call a set small if it is of size less than κ . For $M \prec N$, define

$$A \bar{\downarrow}_M^{(\text{ch})} N \iff \text{for every small } A^- \subseteq A \text{ and } N^- \prec N, \\ \text{there is } B^- \subseteq M \text{ such that } B^- \equiv_{N^-} A^-.$$

One can readily check that $\overset{\text{(ch)}}{\perp}$ satisfies the properties of an independence relation. $\overset{\text{(ch)}}{\perp}$ was first studied in [BG], based on results of [MS90] and [Bon14b], and generalizes the first-order notion of coheir. An alternative name for this notion is $(< \kappa)$ *satisfiability*. Sufficient conditions for this relation to be well-behaved (i.e. to have most of the properties listed above) are given in [BG, Theorem 5.1], reproduced here as Fact 3.16.

Definition 3.8. We define a natural closure for $\overset{\text{(ch)}}{\perp}$:

$$A \overset{\text{(ch)}}{\perp}_M C \iff \text{for every small } A^- \subseteq A \text{ and } C^- \subseteq C, \\ \text{there is } B^- \subseteq M \text{ such that } B^- \equiv_{C^-} A^-.$$

It is straightforward to check that $\overset{\text{(ch)}}{\perp}$ is indeed a closure of $\overset{\text{(ch)}}{\perp}$, but it is not clear at all that this is the *minimal* one. This closure will be useful in the proof of local character (Theorem 6.4) Note that $\overset{\text{(ch)}}{\perp}$ differs from the notion of coheir given in [MS90]; there, types are consistent sets of formulas from a fragment of $L_{\kappa, \kappa}$ for κ strongly compact and the notion there (see [MS90, Definition 4.5]) allows parameters from C and $|M|$.

Definition 3.9 (μ -nonsplitting, [She99]). Let $\mu \geq LS(K)$. For $M \prec N$, we say $A \overset{(\mu\text{-ns})}{\perp}_M N$ if and only if for all $N_1, N_2 \in K_{\leq \mu}$ with $M \prec N_\ell \prec N$, $\ell = 1, 2$, if $f : N_1 \equiv_M N_2$, then there is $g : N_1 \equiv_{AM} N_2$ such that $f \upharpoonright N_1 = g \upharpoonright N_1$.

There is also a definition of nonsplitting that does not depend on a cardinal μ .

Definition 3.10 (Nonsplitting). For $M \prec N$,

$$A \overset{\text{(ns)}}{\perp}_M N \iff A \overset{(\mu\text{-ns})}{\perp}_M N \text{ for all } \mu.$$

An equivalent definition of nonsplitting is given by the following.

Proposition 3.11. $A \overset{\text{(ns)}}{\perp}_M N$ if and only if for all $N_1, N_2 \in K$ with $M \prec N_\ell \prec N$, $\ell = 1, 2$, if $h : N_1 \equiv_M N_2$, then $f : A \equiv_{N_2} h[A]$ for some f with $f \upharpoonright A = h \upharpoonright A$ (equivalently, $\bar{a} \equiv_{N_2} h(\bar{a})$ for all enumerations \bar{a} of A).

The analog statement also holds for μ -nonsplitting.

Proof. Assume $h : N_1 \equiv_M N_2$, and $f : A \equiv_{N_2} h[A]$ is such that $f \upharpoonright A = h \upharpoonright A$. Let $g := f^{-1} \circ h$. Then $g \upharpoonright N_1 = h \upharpoonright N_1$, and g fixes AM . In other words, $g : N_1 \equiv_{AM} N_2$.

N_2 is as needed. Conversely, assume $h : N_1 \equiv_M N_2$. Find $g : N_1 \equiv_{AM} N_2$ such that $h \upharpoonright N_1 = g \upharpoonright N_1$. Then $f := h \circ g^{-1}$ is the desired witness that $A \equiv_{N_2} h[A]$. \square

Using Proposition 3.11 to check base monotonicity, it is easy to see that both $\downarrow^{(ns)}$ and $\downarrow^{(\mu\text{-ns})}$ are independence relations. These notions of splitting in AECs were first explored in [She99], but have seen a wide array of uses; see [SV99, Van06, Van13, GVV, Vasa] for examples. μ -nonsplitting is more common in the literature, but we focus on nonsplitting here. Using tameness, there is a correspondence between the two:

Proposition 3.12. Let $M \prec N$ and $\mu \geq LS(K)$. If K is μ -tame for $|A|$ -length types and $\mu' \in [\mu, \|N\|]$, then

$$A \downarrow_M^{(\mu\text{-ns})} N \implies A \downarrow_M^{(\mu'\text{-ns})} N$$

Proof. We use the equivalence given by Proposition 3.11. Let $\mu' \in [\mu, \|N\|]$, and suppose $A \not\downarrow_M^{(\mu'\text{-ns})} N$. Then there are $N_\ell \in K_{\mu'}$ so $M \prec N_\ell \prec N$ for $\ell = 1, 2$ and $h : N_1 \equiv_M N_2$, but $\bar{a} \not\equiv_{N_2} h(\bar{a})$ for some enumeration \bar{a} of A . By tameness, there is $N_2^- \in K_{\leq \mu}$ so that $\bar{a} \not\equiv_{N_2^-} h(\bar{a})$. Without loss of generality, $M \prec N_2^-$. Let $N_1^- := h^{-1}[N_2^-]$. Then N_1^- and N_2^- witness that $A \not\downarrow_M^{(\mu\text{-ns})} N$. \square

A variant is explicit nonsplitting, which allows the N_i 's to be sets instead of requiring models; this is based on explicit non-strong splitting from [She99, Definition 4.11.2].

Definition 3.13 (Explicit Nonsplitting). For $M \prec N$, we say $A \downarrow_M^{(nes)} N$ if and only if for all $C_1, C_2 \subseteq N$, if $f : C_1 \equiv_M C_2$, then there is $g : C_1 \equiv_{AM} C_2$ such that $f \upharpoonright C_1 = g \upharpoonright C_1$.

From the definition, we see immediately that $\downarrow^{(nes)} \subseteq \downarrow^{(ns)}$. Of course, the corresponding version of Proposition 3.11 also holds for $\downarrow^{(nes)}$, so it is again straightforward to check that $\downarrow^{(nes)}$ is an independence relation. One advantage of using $\downarrow^{(nes)}$ over $\downarrow^{(ns)}$ is that it has a natural closure:

Definition 3.14. We say $A \downarrow_M^{(\overline{nes})} C$ if and only if for all $C_1, C_2 \subseteq C$, if $f : C_1 \equiv_M C_2$, then there is $g : C_1 \equiv_{AM} C_2$ such that $f \upharpoonright C_1 = g \upharpoonright C_1$.

Again, it is not clear this is the minimal closure. We will have no use for this closure, so for most of the paper we will stick with regular nonsplitting.

Nonsplitting will be used mostly as a technical tool to state and prove intermediate lemmas, while coheir will be relevant only in Section 6.

3.3. Properties of coheir and nonsplitting. We now investigate the properties satisfied by coheir and nonsplitting. Here is what holds in general:

Proposition 3.15. Let $\kappa > \text{LS}(K)$.

- (1) $\downarrow^{(\text{ch})}$ and $\downarrow^{(\bar{\text{ch}})}$ have $(C)_\kappa$, and (T) .
- (2) If M is κ -saturated, $\downarrow^{(\text{ch})}$ and $\downarrow^{(\bar{\text{ch}})}$ have $(E_0)_M$.
- (3) $\downarrow^{(\text{ns})}$, $\downarrow^{(\text{nes})}$, and $\downarrow^{(\bar{\text{nes}})}$ have (E_0) .

Proof. Just check the definitions. □

While extension and uniqueness are usually considered very strong assumptions, it is worth noting that nonsplitting satisfies a weak version of them, see [Van06, Theorems I.4.10, I.4.12]. It is also well known that nonsplitting has local character assuming tameness and stability (see e.g. [GV06b, Fact 4.6]). This will not be used. Regarding coheir, the following⁴ appears in [BG] :

Fact 3.16. Let $\kappa > \text{LS}(K)$ be regular. Assume K is fully $(< \kappa)$ -tame, fully $(< \kappa)$ -type short, has no weak κ -order property⁵ and $\downarrow^{(\text{ch})}$ has (E) ⁶.

Then $\downarrow^{(\text{ch})}$ has (U) and (S) .

Moreover, if κ is strongly compact, then the tameness and type-shortness hypotheses hold for free, $\downarrow^{(\text{ch})}$ has (E_1) , and “no weak κ order property” is implied by “ $\exists \lambda > \kappa$ so $I(\lambda, K) < 2^\lambda$.”

As we will see, right transitivity (T_*) can be deduced either from symmetry and (T) (Lemma 5.9) or from uniqueness (Lemma 5.11). Local character will be shown to follow from symmetry (Theorem 6.4).

⁴Since this paper was first submitted, a stronger result has been proven (for example one need not assume (E)). See [Vasc, Theorem 5.15].

⁵See [BG, Definition 4.2].

⁶All the properties mentioned in this Lemma are valid for models of size $\geq \kappa$ only.

4. COMPARING TWO INDEPENDENCE RELATIONS

In this section, we prove the main result of this paper (canonicity of forking), modulo some extra hypotheses that will be eliminated in Section 5. After discussing some preliminary lemmas, we introduce a strengthening of the extension property, (E_+) , which plays a crucial role in the proof. We then prove canonicity using (E_+) (Corollary 4.8). Finally, we show (E_+) follows from some of the more classical properties that we had previously introduced (Corollary 4.13), obtaining the main result of this section (Corollary 4.14). We conclude by giving some examples showing our hypotheses are close to optimal.

For the rest of this section, we fix two independence relations $\overset{(1)}{\perp}$ and $\overset{(2)}{\perp}$. Recall from Definition 3.1 that this means they satisfy (I) , (M) and (B) . We aim to show that if $\overset{(1)}{\perp}$ and $\overset{(2)}{\perp}$ satisfy enough of the properties introduced in Section 3, then $\overset{(1)}{\perp} = \overset{(2)}{\perp}$.

The first easy observation is that given some uniqueness, only one direction is necessary⁷:

Lemma 4.1. Let M be a model. Assume:

- (1) $\overset{(1)}{\perp}_M \subseteq \overset{(2)}{\perp}_M$
- (2) $(E^{(1)})_M, (U^{(2)})_M$

Then $\overset{(1)}{\perp}_M = \overset{(2)}{\perp}_M$.

Proof. Assume $A \overset{(2)}{\perp}_M N$. By $(E^{(1)})_M$, find $A' \equiv_M A$ so that $A' \overset{(1)}{\perp}_M N$. By hypothesis (1), $A' \overset{(2)}{\perp}_M N$. By $(U^{(2)})_M$, $A' \equiv_N A$. By $(I^{(1)})_M$, $A \overset{(1)}{\perp}_M N$. \square

With a similar idea, one can relate an arbitrary independence relation to nonsplitting⁸:

Lemma 4.2. Assume $(U)_M$. Then $\overset{(ns)}{\perp}_M \subseteq \overset{(1)}{\perp}_M$.

⁷Shelah states as an exercise a variation of this lemma in [She09a, Exercise II.6.6.(1)].

⁸Shelah gives a variation of this lemma in [She09a, Claim III.2.20.(1)].

Proof. Assume $A \downarrow_M N$. Let $M \prec N_1, N_2 \prec N$ and $h : N_1 \equiv_M N_2$. By monotonicity, $A \downarrow_M N_\ell$ for $\ell = 1, 2$. By invariance, $h[A] \downarrow_M N_2$. By $(U)_M$, there is $f : A \equiv_{N_2} h[A]$ with $f \upharpoonright A = h \upharpoonright A$. By Proposition 3.11, $A \downarrow_M^{(ns)} N$. \square

A similar result holds for $\downarrow^{(nes)}$, see Lemma 5.6.

The following consequence of invariance will be used repeatedly:

Lemma 4.3. Assume \downarrow satisfies $(E_1)_M$. Assume $A \downarrow_M N$, and $N' \succ N$. Then there is $N'' \equiv_N N'$ such that $A \downarrow_M N''$.

Proof. By $(E_1)_M$, there is $f : A' \equiv_N A$, $A' \downarrow_M N'$. Thus $f : (A', N') \equiv_N (A, f[N'])$, so letting $N'' := f[N']$ and applying invariance, we obtain $A \downarrow_M N''$. \square

Even though we will not use it, we note that an analogous result holds for left extension, see Lemma 5.8.

We now would like to strengthen Lemma 4.3 as follows: suppose we are given A , $M \prec N_0 \prec N$, and assume N is “very big” (e.g. it is $(2^{|A| + \|N_0\|})^+$ -saturated), but does *not* contain A . Can we find $N'_0 \equiv_M N_0$ with $A \downarrow_M N'_0$, and $N'_0 \prec N$?

We give this property a name:

Definition 4.4 (Strong extension). An independence relation \downarrow has (E_+) (strong extension) if for any $M \prec N_0$ and any set A , there is $N \succ N_0$ such that for all $N' \equiv_{N_0} N$, there is $N'_0 \equiv_M N_0$ with $A \downarrow_M N'_0$ and $N'_0 \prec N'$.

Intuitively, (E_+) says that no matter which isomorphic copy N' of N we pick, even if N' does not contain A , N' is so big that we can still find N'_0 inside N' with the right property. This is stronger than (E) in the following sense:

Proposition 4.5. If \downarrow has (E_+) , \downarrow has (E_0) . If in addition \downarrow has (T_*) , then \downarrow has (E_1) . Thus if \downarrow has (E_+) and (T_*) , it has (E) .

Proof. Use monotonicity and Remark 3.3. \square

Remark 4.6. Example 4.15 shows (E_+) does *not* follow from (E) .

Strong extension allows us to prove canonicity:

Lemma 4.7. Assume $(E_1^{(1)})_M, (E_+^{(2)})_M$. Assume also that $\downarrow_M \subseteq \downarrow_M^{(1)}$ and $\downarrow_M \subseteq \downarrow_M^{(ns)}$.

Then $\downarrow_M^{(1)} \subseteq \downarrow_M^{(2)}$.

Proof. Assume $A \downarrow_M^{(1)} N_0$. We show $A \downarrow_M^{(2)} N_0$. Fix $N \succ N_0$ as described by $(E_+^{(2)})_M$.

By Lemma 4.3, we can find $N' \equiv_{N_0} N$ such that $A \downarrow_M^{(1)} N'$. By definition of N , one can pick $N'_0 \equiv_M N_0$ with $N'_0 \prec N'$ and $A \downarrow_M^{(2)} N'_0$.

We have $A \downarrow_M^{(\text{ns})} N'$, $M \prec N'_0$, $N_0 \prec N'$, and $N'_0 \equiv_M N_0$, so by definition of nonsplitting, $N'_0 \equiv_{AM} N_0$. By invariance, $A \downarrow_M^{(2)} N_0$, as needed. \square

Corollary 4.8 (Canonicity of forking from strong extension). Assume:

- $(U^{(1)})_M, (E^{(1)})_M$.
- $(U^{(2)})_M, (E_+^{(2)})_M$.

Then $\downarrow_M^{(1)} = \downarrow_M^{(2)}$.

Proof. By Lemma 4.1, it is enough to see $\downarrow_M^{(1)} \subseteq \downarrow_M^{(2)}$. By Lemma 4.2, $\downarrow_M^{(1)} \subseteq \downarrow_M^{(\text{ns})}$. The result now follows from Lemma 4.7. \square

We now proceed to show that (E_+) follows from (E) , (T_*) , (S) and (L) . We will use the following important concept:

Definition 4.9 (Independent sequence). Let I be a linearly ordered set. A sequence of sets $(A_i)_{i \in I}$ is *independent* over a model M if there is a strictly increasing continuous chain of models $(N_i)_{i \in I}$ such that for all $i \in I$:

- (1) $M \cup \bigcup_{j < i} A_j \subseteq N_i$ and $N_0 = M$.
- (2) $A_i \downarrow_M N_i$.

This generalizes the notion of independent sequence from the first-order case. The most natural definition would only require $A_i \bar{\downarrow}_M \bigcup_{j < i} A_j$ (for some closure $\bar{\downarrow}$ of \downarrow) but it turns out it is convenient to have a sequence of models $(N_i)_{i \in I}$ witnessing the independence in a uniform way.

We note that very similar definitions appear already in the literature. See [JS12, Definition 3.2], [She09a, Section III.5], or [She09b, Definition V.D.3.15].

Just like in the first-order case, the extension property allows us to build independent sequences:

Lemma 4.10 (Existence of independent sequences). Assume $(E)_M$. Let A be a set, and let δ be an ordinal. Then there is a sequence $(A_i)_{i < \delta}$ independent over M so that $A_i \equiv_M A$ for all $i < \delta$, and $A_0 = A$.

Proof. Define the $(A_i)_{i < \delta}$ and the $(N_i)_{i < \delta}$ witnessing the independence of the sequence by induction on $i < \delta$. Take $N_0 = M$ and $A_0 = A$. Assume inductively $(A_j)_{j < i}, (N_j)_{j < i}$ have been defined. If i is a limit, let $N_i := \bigcup_{j < i} N_j$. If i is a successor, let N_i be any model containing $M \cup \bigcup_{j < i} (A_j \cup N_j)$ and strictly extending the previous N_j 's. By $(E)_M$, there is $A_i \equiv_M A$ such that $A_i \perp_M N_i$. Thus $(A_i)_{i < \delta}$ is as desired. \square

The next result is key to the proof of (E_+) . It is adapted from [Bal88, Theorem II.2.18].

Lemma 4.11. Assume \perp has $(S), (T_*)_M, (L)$. Let A be a set, and let $\mu := \kappa_{|A|}(\perp)$. Then whenever $(M_i)_{i < \mu}$ is an independent sequence over M with $M \prec M_i$ for all i , there is $i < \mu$ with $A \perp_M M_i$.

Proof. Let $(N_i)_{i < \mu}$ witness independence of the M_i 's. Let $N_\mu := \bigcup_{i < \mu} N_i$. By definition of μ , there is $i < \mu$ so that $A \perp_{N_i} N_\mu$. By (S) , there is a model N_A with $N_i \prec N_A$, $A \subseteq N_A$, and $N_\mu \perp_{N_i} N_A$. By (M) , $M_i \perp_{N_i} N_A$. Since the M_i 's are independent, we also have $M_i \perp_M N_i$. By $(T_*)_M$, $M_i \perp_M N_A$. By (S) (recall that $M \prec M_i$), $N_A \perp_M M_i$. By (M) , $A \perp_M M_i$, as desired. \square

Remark 4.12. The same proof works if we replace \perp by its minimal closure $\overline{\perp}$, and $(M_i)_{i < \mu}$ by an arbitrary sequence $(B_i)_{i < \mu}$ independent over M .

Corollary 4.13. Assume $(E)_M, (S), (T_*)_M$, and (L) . Then $(E_+)_M$.

Proof. Fix A and $N_0 \succ M$. Let $\mu := \kappa_{|A|}(\perp)$. By Lemma 4.10, there is a sequence $(M_i)_{i < \mu}$ independent over M such that $M_i \equiv_M N_0$ for all $i < \mu$, and $M_0 = N_0$. Let $(N'_i)_{i < \mu}$ witness independence of the M_i 's. We claim $N := \bigcup_{i < \mu} N'_i$ is as required. By construction, $N_0 = M_0 \prec N$.

Now let $f : N \equiv_{N_0} N'$. Let $M'_i := f[M_i]$. Invariance implies $(M'_i)_{i < \mu}$ is an independent sequence over M inside N' , with $M'_i \equiv_M N_0$ for all $i < \mu$. By Lemma 4.11, there is $i < \mu$ so that $A \perp_M M'_i$, so $N'_0 := M'_i$ is exactly as needed. \square

Corollary 4.14. Assume:

- $(E^{(1)})_M, (U^{(1)})_M$.
- $(E^{(2)})_M, (U^{(2)})_M, (L^{(2)}), (S^{(2)}), (T_*^{(2)})_M$.

Then $\underset{M}{\downarrow}^{(1)} = \underset{M}{\downarrow}^{(2)}$.

Proof. Combine Corollaries 4.8 and 4.13. \square

We will see (Corollary 5.18) that (S) and (T_*) follow from (E) , (U) , and (L) . We now argue that the other hypotheses are necessary. The following example (versions of which appears at various places in the literature, e.g. [She09a, Example II.6.4], [Adl09a, Example 6.6]) shows we cannot remove the local character assumption from Corollary 4.14. In particular, (E_+) does not follow from (E) and (U) alone. The example also shows the AxFr₃ framework (see [She09b, Definition V.B.1.9]) is not canonical.

Example 4.15. Let T_{ind} be the first-order theory of the random graph, and let K be the class of models of T_{ind} , ordered by first-order elementary substructure. Define

- $A \underset{M}{\downarrow}^{(1)} N$ iff $A \cap N \subseteq M$, and there are no edges between $A \setminus M$ and $N \setminus M$.
- $A \underset{M}{\downarrow}^{(2)} N$ iff $A \cap N \subseteq M$, and all the possible cross edges between $A \setminus M$ and $N \setminus M$ are present.

It is routine to check that both $\underset{M}{\downarrow}^{(1)}$ and $\underset{M}{\downarrow}^{(2)}$ are independence relations with (E) , (U) , (S) , (T) , (T_*) , $(C)_{\aleph_0}$. Yet $\underset{M}{\downarrow}^{(1)} \neq \underset{M}{\downarrow}^{(2)}$, so one knows from Corollary 4.14 (or from first-order stability theory) that K can have *no* independence relation which in addition has (L) or (E_+) .

Of course, T_{ind} is simple, so first-order nonforking will actually have (E_+) , local character, transitivity and symmetry (but not uniqueness).

A concrete reason (E_+) does not hold e.g. for $\underset{M}{\downarrow}^{(1)}$ is that given $M \prec N_0$ one can pick $a \notin N_0$ such that there is an edge from a to any element of N_0 . Then for any $N \succ N_0$, one can again pick $N' \equiv_{N_0} N$, disjoint from $\{a\} \cup (N \setminus N_0)$ such that there is an edge from a to any element of N' . Then $a \underset{M}{\downarrow}^{(1)} N'_0$ for $N'_0 \prec N$ implies $N'_0 = M$. Local character fails for a similar reason.

Example 4.16. It is also easy to see that $(E^{(2)})$ and $(U^{(2)})$ are necessary in Corollary 4.14. Assume $\underset{M}{\downarrow}^{(1)}$ has (E) , (U) , (S) , (T_*) , and (L) . Then the independence

relation $\overset{(2)}{\perp}$ defined by $A \overset{(2)}{\perp}_M N$ for all A and $M \prec N$ satisfies $(E), (S), (T_*), (L)$, but not (U) , so is distinct from $\overset{(1)}{\perp}$.

Similarly define $A \overset{(2)}{\perp}_M N$ if and only if $M \prec N$ and either both $A \overset{(1)}{\perp}_M N$ and $\|M\| \geq \text{LS}(K)^+$, or $M = N$. Then $\overset{(2)}{\perp}$ has $(E_0), (U), (S), (T_*)$ and (L) , but does not have $(E_1)_M$ if M is a model of size $\text{LS}(K)$. This last example was adapted from [Adl09a, Example 6.4].

Remark 4.17. After the initial submission of this paper, it was shown in [Vasb, Lemma 9.1] that (E) can be removed from the hypotheses of Corollary 4.14 (but one has to replace it by $(C)_\kappa$) if one only wants the independence relations to agree over sufficiently saturated models.

5. RELATIONSHIP BETWEEN VARIOUS PROPERTIES

In this section, we investigate some of relations between the properties introduced earlier. We first discuss the interaction between properties of an independence relation and properties of its closures, and show how to obtain transitivity from various other properties. We then show how to obtain symmetry from existence, extension, uniqueness, and local character (Corollary 5.18). This second part has a stability-theoretic flavor and most of it does not depend on the first part.

Most of the material in the first part of this section is not used in the rest of the paper, but the concept of closure (Definition 3.4) felt unmotivated without it. Our investigation remains far from exhaustive, and leaves a lot of room for further work.

5.1. Properties of the minimal closure. Recall the notion of closure of an independence relation (Definition 3.4). We would like to know when we can transfer properties from an independence relation to its closures and vice-versa.

For an arbitrary closure, we can say little:

Lemma 5.1. Let $\bar{\perp}$ be a closure of \perp . Then:

- (1) A property in the following list holds for \perp if and only if it holds for $\bar{\perp}$: $(T_*)_M, (E_0)_M, (L)$.
- (2) If a property in the following list holds for $\bar{\perp}$, then it holds for \perp : $(C)_\kappa, (T)_M, (E_1)_M, (U)_M$.

Proof.

- (1) Because those properties have the same definition for \perp and $\bar{\perp}$.
- (2) Straightforward from the definitions.

□

The minimal closure is more interesting. We start by generalizing Lemma 4.3:

Lemma 5.2. Assume \perp satisfies $(E_1)_M$. Let $\bar{\perp}$ be the minimal closure of \perp . Assume $A \bar{\perp}_M C$, and let B be an arbitrary set. Then there is $B' \equiv_{MC} B$ such that $A \bar{\perp}_M B'$.

Proof. Let N be a model containing C and M such that $A \perp_M N$. Let N' be a model containing NB . By Lemma 4.3, there is $N'' \equiv_N N'$ such that $A \perp_M N''$. Now use monotonicity to get the result. □

The next lemma tells us that the minimal closure is the only one that will keep the extension property:

Lemma 5.3. Let $\bar{\perp}$ be a closure of \perp and let $\bar{\perp}$ be the minimal closure of \perp . Assume \perp has $(E_1)_M$. Then $\bar{\perp} = \bar{\perp}$ if and only if $\bar{\perp}$ has $(E_1)_M$.

Proof. Assume first $\bar{\perp} = \bar{\perp}$. Let $C \subseteq C'$, and assume $A \bar{\perp}_M C$. Then by definition of the minimal closure, there exists $N \succ M$ containing C such that $A \perp_M N$. Let N' be a model containing N and C' . By $(E_1)_M$ for \perp , there is $A' \equiv_N A$ so that $A' \perp_M N'$. By monotonicity, $A' \bar{\perp}_M C'$, and since N contains C , $A' \bar{\perp}_M C$.

Conversely, assume $\bar{\perp}$ has $(E_1)_M$. We know already that $\bar{\perp} \subseteq \bar{\perp}$, so assume $A \bar{\perp}_M C$. Let N be a model containing M and C . By Lemma 5.2, there is $N' \equiv_{MC} N$ so that $A \bar{\perp}_M N'$, so $A \bar{\perp}_M C$, as needed. □

Lemma 5.4. Let $\bar{\perp}$ be the *minimal* closure of \perp . Then

- (1) $(E)_M$ holds for \perp if and only if it holds for $\bar{\perp}$.
- (2) $(S)_M$ holds for \perp if and only if it holds for $\bar{\perp}$.

- (3) If \perp has $(E)_M$, then it has $(U)_M$ if and only if $\bar{\perp}$ does.
(4) If \perp has (E) , then it has (T) if and only if $\bar{\perp}$ does⁹.

Proof.

- (1) By Lemmas 5.1 and 5.3.
(2) Straightforward from the definition of symmetry and monotonicity.
(3) One direction holds by Lemma 5.1. For the other direction, assume $\bar{\perp}$ has $(E_1)_M$ and \perp has $(U)_M$. Assume $A \bar{\perp}_M C$ and $A' \bar{\perp}_M C$, with $f : A \equiv_M A'$. Let N be a model containing MC such that $A \perp_M N$. By extension again, find $h : A' \equiv_{MC} A''$ such that $A'' \perp_M N$. We know $h' := h \circ f : A \equiv_M A''$, so by uniqueness, there is $h'' : A \equiv_{N'} A''$, and $h'' \upharpoonright A = h' \upharpoonright A = (h \circ f) \upharpoonright A$, so $f \upharpoonright A = (h^{-1} \circ h'') \upharpoonright A$. Therefore $g := h^{-1} \circ h''$ is the desired witness that $A \equiv_{MC} A'$.
(4) Let $M_0 \prec M_1 \prec M_2$, and assume $M_1 \bar{\perp}_{M_0} C$, $M_2 \bar{\perp}_{M_1} C$. Let N be an extension of M_1 containing C such that $M_2 \bar{\perp}_{M_1} N$. Let χ be a big cardinal, so that (V_χ, \in) reflects enough set theory and contains NM_2 . Let N' be what V_χ believes is the monster model. By Lemma 4.3, there is $f : N'' \equiv_N N'$ such that $M_2 \perp_{M_1} N''$. Notice that $CM_1 \subseteq N \subseteq N'$, so since we took χ big enough, we can apply the definition of the minimal closure *inside* V_χ to get $N'_0 \prec N'$ containing M_0 and C so that $M_1 \perp_{M_0} N'_0$. Let $N_0 := f[N'_0]$. By invariance, $M_1 \perp_{M_0} N_0$, and $N_0 \prec N''$, so by monotonicity, $M_2 \perp_{M_1} N_0$, so by $(T)_{M_0}$ for \perp , $M_2 \perp_{M_0} N_0$. By monotonicity again, $M_2 \bar{\perp}_{M_0} C$.

□

The following remains to be investigated:

Question 5.5. Let $\bar{\perp}$ be the minimal closure of \perp . Under what conditions does $(C)_\kappa$ for \perp imply $(C)_\kappa$ for $\bar{\perp}$?

⁹More precisely, if \perp has $(E)_{M_1}$, and for $M_0 \prec M_1 \prec M_2$, we have that $M_2 \perp_{M_1} N$, $M_1 \perp_{M_0} N$ implies $M_2 \perp_{M_0} N$, then $M_2 \bar{\perp}_{M_1} C$, $M_1 \bar{\perp}_{M_0} C$ implies $M_2 \bar{\perp}_{M_0} C$.

We can use Lemma 5.4 to prove a variation on Lemma 4.2.

Lemma 5.6. Assume \perp has $(E)_M$ and $(U)_M$. Then $\perp \subseteq \overset{\text{(nes)}}{\perp}_M$.

Proof. Let $\bar{\perp}$ be the minimal closure of \perp . By Lemma 5.4, $\bar{\perp}$ has $(E)_M$ and $(U)_M$.

Assume $A \underset{M}{\perp} N$. Let $C_1, C_2 \subseteq N$, and $h : C_1 \equiv_M C_2$. By monotonicity, $A \underset{M}{\bar{\perp}} C_\ell$ for $\ell = 1, 2$. By invariance, $h[A] \underset{M}{\bar{\perp}} C_2$. By $(U)_M$, there is $f : A \equiv_{MC_2} h[A]$ with $f \upharpoonright A = h \upharpoonright A$. By (the proof of) Proposition 3.11, $A \overset{\text{(nes)}}{\underset{M}{\perp}} N$. \square

Question 5.7. Is the $(E)_M$ hypothesis necessary?

We can also obtain a left version of Lemma 4.3:

Lemma 5.8. Let $\bar{\perp}$ be a closure of \perp . Assume \perp has $(E)_N$, and $\bar{\perp}$ has $(T)_{M_1}$. Suppose that $N \underset{M_1}{\perp} M_2$, with $N \succ M_1$. Then for all $N' \succ N$, there exists $N'' \equiv_N N'$ such that $N'' \underset{M_1}{\perp} M_2$.

In particular, this holds if \perp has (E) and (T) .

Proof. The last line follows from part (4) of Lemma 5.4 by taking $\bar{\perp}$ to be the minimal closure of \perp .

To see the rest, let N_3 be a model containing M_2N . By $(E)_N$, there is $N'' \equiv_N N'$ such that $N'' \underset{N}{\perp} N_3$. Since $M_2 \subseteq N_3$, $N'' \underset{N}{\bar{\perp}} M_2$. By hypothesis, $N \underset{M_1}{\bar{\perp}} M_2$. So since $\bar{\perp}$ has $(T)_{M_1}$, $N'' \underset{M_1}{\bar{\perp}} M_2$. Since $M_2 \succ M_1$, $N'' \underset{M_1}{\perp} M_2$. \square

Finally, we can also use symmetry to translate between the transitivity properties:

Lemma 5.9. Assume \perp has (S) . Then:

- (1) If \perp has $(T_*)_{M_0}$, then \perp has $(T)_{M_0}$.
- (2) If \perp has $(T)_{M_0}$ and (E) , then it has $(T_*)_{M_0}$.

Proof. Let $M_0 \prec M_1 \prec M_2$. Let $\bar{\perp}$ be the minimal closure of \perp . By Lemma 5.4, $\bar{\perp}$ has (S) .

- (1) By Lemma 5.1, $\bar{\perp}$ has $(T_*)_{M_0}$. Now use symmetry.

(2) By part (4) of Lemma 5.4, $\bar{\downarrow}$ has $(T)_{M_0}$. Now use symmetry. □

This gives us one way to obtain right transitivity for coheir:

Corollary 5.10. Assume \downarrow has (S) and (E) . Then \downarrow has (T_*) .

Proof. By Proposition 3.15, \downarrow has (T) . Apply Lemma 5.9. □

Another way to obtain right transitivity from other properties appears in [She09a, Claim II.2.18]:

Lemma 5.11. Assume \downarrow has $(E_1)_M$ and (U) . Then \downarrow has $(T_*)_M$.

Proof. Let $M_0 \prec M_1 \prec M_2$, and assume $A \downarrow_{M_0} M_1$ and $A \downarrow_{M_1} M_2$. By $(E_1)_M$, there exists $A' \equiv_{M_1} A$ such that $A' \downarrow_{M_0} M_2$. By base monotonicity, $A' \downarrow_{M_1} M_2$. By uniqueness, $A \equiv_{M_2} A'$. By invariance, $A \downarrow_{M_0} M_2$. □

5.2. Getting symmetry. We prove that symmetry follows from (E) , uniqueness and local character and deduce the main theorem of this paper (Corollary 5.19). We start by assuming some stability. The following is a strengthening of unstability that is sometimes more convenient to work with:

Definition 5.12. Let α and λ be cardinals. K has the α -order property of length λ if there is a sequence $(\bar{a}_i)_{i < \lambda}$ of tuples, with $\ell(\bar{a}_i) = \alpha$, so that for any $i_0 < j_0 < \lambda$ and $i_1 < j_1 < \lambda$, $\bar{a}_{i_0} \bar{a}_{j_0} \not\equiv \bar{a}_{j_1} \bar{a}_{i_1}$.

K has the α -order property if it has the α -order property of all lengths.

K has the order property if it has the α -order property for some cardinal α .

This is a variation on the order property defined in [She99] taken from [GV06b]. It is stronger than unstability:

Fact 5.13. Let α be a cardinal. If K has the α -order property, then K is α -unstable.

Proof sketch. This is [She99, Claim 4.7.2]. Shelah's proof is "Straight.", so we elaborate a little.

Let $\lambda \geq \text{LS}(K)$. We show K is α -unstable in λ . Let $I \subseteq \hat{I}$ be linear orderings such that $\|I\| \leq \lambda$, $\|\hat{I}\| > \lambda$, and I is dense in \hat{I} . Combining Shelah's presentation

theorem with Morley's method, we can get a sequence $\hat{\mathbf{I}} := \langle \bar{a}_i \mid i \in \hat{I} \rangle$ with $\ell(\bar{a}_i) = \alpha$ and $i_0 < j_0, i_1 < j_1$ implies $\bar{a}_{i_0}\bar{a}_{j_0} \not\equiv \bar{a}_{j_1}\bar{a}_{i_1}$. Let $\mathbf{I} := \langle \bar{a}_i \mid i \in I \rangle$.

Now for any $i < j$ in \hat{I} , $\bar{a}_i \not\equiv_{\mathbf{I}} \bar{a}_j$. Indeed, pick $i < k < j$ with $k \in I$. Then $\bar{a}_i\bar{a}_k \not\equiv \bar{a}_j\bar{a}_k$ by construction, so $\bar{a}_i \not\equiv_{\bar{a}_k} \bar{a}_j$. This completes the proof that K is α -unstable in λ . \square

We are now ready to prove symmetry. The argument is similar to [She90, Theorem III.4.13] or [She75, Theorem 5.1].

Theorem 5.14 (Symmetry). Assume \downarrow has $(E)_M$ and $\downarrow \subseteq \downarrow \stackrel{(\text{nes})}{\subseteq} \downarrow$. Assume in addition that K does not have the order property. Then \downarrow has $(S)_M$.

Proof. Let $\bar{\downarrow}$ be the minimal closure of \downarrow . Recall that by Lemma 5.4, \downarrow has $(S)_M$ if and only if $\bar{\downarrow}$ has $(S)_M$.

Assume for a contradiction \downarrow does not have $(S)_M$. Pick A and $M \prec N$ such that $A \downarrow_M N$, but $N \not\downarrow_M A$. Let λ be an arbitrary uncountable cardinal. We will show that K has the $(\|N\| + |A|)$ -order property of length λ . This will contradict the assumption that K does not have the order property.

We will build increasing continuous $\langle M_\alpha \in K : \alpha < \lambda \rangle$, and $\langle A_\alpha, M'_\alpha, N_\alpha : \alpha < \lambda \rangle$ by induction so

- (1) $M_0 \succ N$ and $A \subseteq |M_0|$.
- (2) $N_\alpha \equiv_M N$ and $N_\alpha \prec M'_\alpha$.
- (3) $A_\alpha \equiv_N A$ and $A_\alpha \subseteq M_{\alpha+1}$.
- (4) $M_\alpha \prec M'_\alpha \prec M_{\alpha+1}$.
- (5) $N_\alpha \downarrow_M M_\alpha$ and $A_\alpha \downarrow_M M'_\alpha$.

This is possible. Let M_0 be any model containing AN . At α limits, let $M_\alpha := \bigcup_{\beta < \alpha} M_\beta$. Now assume inductively that M_β has been defined for $\beta \leq \alpha$, and $A_\beta, N_\beta, M'_\beta$ have been defined for $\beta < \alpha$. Use $(E)_M$ to find $N_\alpha \equiv_M N$ with $N_\alpha \downarrow_M M_\alpha$.

Now pick $M'_\alpha \geq M_\alpha$ containing N_α . Now, by $(E)_M$ again, find $A_\alpha \equiv_N A$ with $A_\alpha \downarrow_M M'_\alpha$. Pick $M_{\alpha+1} \succ M_\alpha$ containing A_α and M'_α .

This is enough. We show that for $\alpha, \beta < \lambda$:

- (1) If $\beta < \alpha$, $(A, N) \not\equiv_M (A_\beta, N_\alpha)$.
- (2) If $\beta \geq \alpha$, $(A, N) \equiv_M (A_\beta, N_\alpha)$.

For (1), suppose $\beta < \alpha$. Since $A \subseteq M \prec M_\alpha$, we have $N_\alpha \downarrow_M^{\bar{}} A$. Then we can use

the invariance of $\bar{\downarrow}$ and the assumption of no symmetry to conclude $(A, N_\alpha) \not\equiv_M (A, N)$. On the other hand, we know that $N_\alpha \downarrow_M^{(\text{nes})} M_\alpha$. Since $A, A_\beta \subseteq M_\alpha$ and $A \equiv_M A_\beta$, we must have $(A, N_\alpha) \equiv_M (A_\beta, N_\alpha)$. Thus $(A, N) \not\equiv_M (A_\beta, N_\alpha)$.

To see (2), suppose $\beta \geq \alpha$ and recall that $(A, N) \equiv_M (A_\beta, N)$. We also have that $A_\beta \downarrow_M^{(\text{nes})} M'_\beta$. $N \equiv_M N_\alpha$ and $N, N_\alpha \subseteq M'_\beta$, the definition of non explicit splitting implies that $(A_\beta, N) \equiv_M (A_\beta, N_\alpha)$. This gives us that $(A, N) \equiv_M (A_\beta, N_\alpha)$ as desired.

□

Remark 5.15. The same proof can be used to obtain symmetry in the good frame framework. This is used in the construction of a good frame of [Vasa].

Corollary 5.16. Assume K does not have the order property. Assume \downarrow has $(E)_M$ and $(U)_M$. Then \downarrow has $(S)_M$.

Proof. By Lemma 5.6, $\downarrow \subseteq_M^{(\text{nes})} \downarrow_M$. Now apply Theorem 5.14. □

If in addition we assume local character, we obtain the “no order property” hypothesis:

Lemma 5.17. Assume \downarrow has (U) and (L) (or just $\kappa_1(\downarrow) < \infty$). Then K is α -stable for all α . In particular, it does not have the order property.

Proof. That α -stability implies no α -order property is the contrapositive of Fact 5.13. Now, assume (U) and let $\mu := \kappa_1(\downarrow) < \infty$. Fix a cardinal $\alpha \geq 1$. We want to see K is α -stable. By Remark 2.10, we can assume without loss of generality $\alpha \geq \mu + \text{LS}(K)$.

Let $\lambda := \beth_{\alpha^+}$. Then:

- (1) λ is strong limit.
- (2) $\text{cf}(\lambda) = \alpha^+ > \mu + \text{LS}(K)$.
- (3) $\lambda^\alpha = \sup_{\gamma < \lambda} \gamma^\alpha = \lambda$.

We claim that K is α -stable in λ . By Fact 2.11, it is enough to see it is 1-stable in λ . Suppose not. Then there exists $M \in K_\lambda$, and $\{a_i\}_{i < \lambda^+}$ such that $i < j$ implies $a_i \not\equiv_M a_j$. Let $(M_i)_{i < \lambda}$ be increasing continuous such that $M = \bigcup_{i < \lambda} M_i$ and $\|M_i\| < \lambda$. By definition of μ , for each $i < \lambda^+$, there exists $k_i < \lambda$ such that $a_i \downarrow_{M_{k_i}} M$. By the pigeonhole principle, we can shrink $\{a_i\}_{i < \lambda^+}$ to assume without

loss of generality that $k_i = k_0$ for all $i < \lambda^+$. Since there are at most $2^{\|M_{k_0}\|} < \lambda$ many types over M_{k_0} , there exists $i < j < \lambda^+$ such that $a_i \equiv_{M_{k_0}} a_j$. By uniqueness, $a_i \equiv_M a_j$, a contradiction. \square

Corollary 5.18. Assume \perp has $(E)_M$, (U) and (L) (or just $\kappa_1(\perp) < \infty$). Then \perp has $(S)_M$ and $(T^*)_M$.

Proof. Lemma 5.11 gives $(T^*)_M$. Combine Lemma 5.17 and Corollary 5.16 to obtain $(S)_M$. \square

Thus we obtain another version of the canonicity theorem:

Corollary 5.19 (Canonicity of forking). Let $\perp^{(1)}$ and $\perp^{(2)}$ be independence relations. Assume:

- $(E^{(1)})_M, (U^{(1)})_M$.
- $(E^{(2)})_M, (U^{(2)}), (L^{(2)})$.

Then $\perp^{(1)} = \perp^{(2)}$.

In particular, there can be at most one independence relation satisfying existence, extension, uniqueness, and local character.

Proof. Combine Corollaries 4.14 and 5.18. \square

6. APPLICATIONS

6.1. Canonicity of coheir. Fix a regular $\kappa > \text{LS}(K)$. Below, when we say coheir has a given property, we mean that it has that property for base models in $K_{\geq \kappa}$.

We are almost ready to show that coheir is canonical, but we first need to show it has local character. We will use the following strengthening that deals with subsets instead of chains of models:

Definition 6.1. Let \perp be an independence relation. For α a cardinal, let $\bar{\kappa}_\alpha = \bar{\kappa}_\alpha(\perp)$ be the smallest cardinal such that for all N , and all A with $|A| = \alpha$, there exists $M \leq N$ with $\|M\| < \bar{\kappa}_\alpha$ and $A \perp M$. $\bar{\kappa}_\alpha = \infty$ if there is no such cardinal.

Remark 6.2. For all α , $\kappa_\alpha(\perp) \leq \bar{\kappa}_\alpha(\perp)^+$. Thus $\bar{\kappa}_\alpha(\perp) < \infty$ implies $\kappa_\alpha(\perp) < \infty$.

Remark 6.3. The converse is also true: under some reasonable hypotheses, $\kappa_\alpha(\perp) < \infty$ implies $\bar{\kappa}_\alpha(\perp) < \infty$. This appears as [BV, Proposition 2.15] (circulated after the initial submission of this paper).

Theorem 6.4 (Local character for coheir). Assume \perp has (S) . Then $\bar{\kappa}_\alpha(\perp) \leq ((\alpha + 2)^{< \kappa})^+$. In particular, \perp has (L) .

The proof is similar to that of [Adl09b, Theorem 1.6]¹⁰. The key is that $\downarrow^{(\bar{\text{ch}})}$ always satisfies a dual to local character:

Lemma 6.5. Let N, C be given. Then there is $M \leq N$, $\|M\| \leq (|C| + 2)^{<\kappa} + LS(K)$ such that $N \downarrow_M^{(\bar{\text{ch}})} C$.

Proof sketch. For each of the $|C|^{<\kappa}$ small subsets of C , look at the $\leq 2^{<\kappa}$ small types over that set (realized in N), and collect a realization of each in a set $A \subseteq |N|$. Then pick $M \prec N$ to contain A and be of the appropriate size. \square

We will also use the following application of the fact $\downarrow^{(\bar{\text{ch}})}$ has $(C)_\kappa$ and a strong form of base monotonicity.

Lemma 6.6. Let λ be such that $\text{cf } \lambda \geq \kappa$. Let $(A_i)_{i < \lambda}$, $(M_i)_{i < \lambda}$, $(C_i)_{i < \lambda}$ be (not necessarily strictly) increasing chains. Assume $A_i \downarrow_{M_i}^{(\bar{\text{ch}})} C_i$ for all $i < \lambda$. Let $A_\lambda := \bigcup_{i < \lambda} A_i$, and define M_λ, C_λ similarly. Then $A_\lambda \downarrow_{M_\lambda}^{(\bar{\text{ch}})} C_\lambda$.

Proof. From the definition of $\downarrow^{(\bar{\text{ch}})}$, we see that for all $i < \lambda$, $A_i \downarrow_{M_\lambda}^{(\bar{\text{ch}})} C_i$. Now use the fact that $\downarrow^{(\bar{\text{ch}})}$ has $(C)_\kappa$ (Proposition 3.15). \square

Proof of Theorem 6.4. Fix α , and let A and N be given with $|A| = \alpha$. Let $\mu := (|A| + 2)^{<\kappa}$. Inductively build $(M_i)_{i \leq \mu}, (N_i)_{i \leq \mu}$ increasing continuous such that for all $i < \mu$:

- (1) $A \subseteq N_i$.
- (2) $M_i \prec N$, $\|M_i\| \leq \mu$.
- (3) $M_i \prec N_{i+1}$.
- (4) $N \downarrow_{M_{i+1}}^{(\bar{\text{ch}})} N_{i+1}$.

This is enough: By König's lemma, $\text{cf } \mu \geq \kappa$ so by Lemma 6.6, $N \downarrow_{M_\mu}^{(\bar{\text{ch}})} N_\mu$. Moreover, by (2), (3) and the chain axioms, $M_\mu \prec N_\mu, N$, and by (2), $\|M_\mu\| \leq \mu$. Thus

¹⁰After proving the result, we noticed that a similar argument also appears in the proof of $(B)_\mu$ in [MS90, Proposition 4.8].

$N \underset{M_\mu}{\overset{\text{(ch)}}{\downarrow}} N_\mu$, and one can apply (S) to get $N_\mu \underset{M_\mu}{\overset{\text{(ch)}}{\downarrow}} N$. By monotonicity, $A \underset{M_\mu}{\overset{\text{(ch)}}{\downarrow}} N$, exactly as needed.

This is possible: Pick any $A \subseteq N_0$ with $\|N_0\| \leq \mu$ (this is possible since $\mu \geq |A| + \kappa > \text{LS}(K)$). Now, given i non-limit, $(N_j)_{j \leq i}$ and $(M_j)_{j < i}$, use Lemma 6.5 to find $M_i \prec N$, $\|M_i\| \leq (\|N_i\|)^{<\kappa} \leq \mu$, such that $N \underset{M_i}{\overset{\text{(ch)}}{\downarrow}} N_i$. Then pick any N_{i+1} extending both M_i and N_i , with $\|N_{i+1}\| \leq \mu$. □

We finally have all the machinery to prove:

Theorem 6.7 (Canonicity of coheir). Assume K is fully $(< \kappa)$ -tame, fully $(< \kappa)$ -type short, and has no weak κ -order property¹¹.

Assume $\underset{\downarrow}{\overset{\text{(ch)}}{\downarrow}}$ has (E). Then:

- (1) $\underset{\downarrow}{\overset{\text{(ch)}}{\downarrow}}$ has $(C)_\kappa$, (T), (T_*) , (S), (U), and (L).
- (2) Any independence relation satisfying (E) and (U) must be $\underset{\downarrow}{\overset{\text{(ch)}}{\downarrow}}$ (for base models in $K_{\geq \kappa}$).

Proof. By Proposition 3.15, $\underset{\downarrow}{\overset{\text{(ch)}}{\downarrow}}$ has $(C)_\kappa$ and (T). By Fact 3.16, $\underset{\downarrow}{\overset{\text{(ch)}}{\downarrow}}$ has (U) and (S). By Corollary 5.10 (or Lemma 5.11), $\underset{\downarrow}{\overset{\text{(ch)}}{\downarrow}}$ also has (T_*) . By Theorem 6.4, $\underset{\downarrow}{\overset{\text{(ch)}}{\downarrow}}$ has (L). This takes care of (1). (2) follows from (1) and Corollary 4.14. □

Corollary 6.8 (Canonicity of coheir, assuming a strongly compact). Assume κ is strongly compact, all models in $K_{\geq \kappa}$ are κ -saturated, and there exists $\lambda > \kappa$ such that $I(\lambda, K) < 2^\lambda$. Then:

- (1) $\underset{\downarrow}{\overset{\text{(ch)}}{\downarrow}}$ has (E), $(C)_\kappa$, (T), (T_*) , (S), (U), and (L).
- (2) Any independence relation satisfying (E) and (U) must be $\underset{\downarrow}{\overset{\text{(ch)}}{\downarrow}}$ (for base models in $K_{\geq \kappa}$).

Proof. By Proposition 3.15, $\underset{\downarrow}{\overset{\text{(ch)}}{\downarrow}}$ has (E_0) . Thus by the moreover part of Fact 3.16, $\underset{\downarrow}{\overset{\text{(ch)}}{\downarrow}}$ has (E). Now apply Theorem 6.7. □

¹¹See [BG, Definition 4.2].

6.2. Canonicity of good frames. As has already been noted, the framework AxFri_3 defined in [She09b, Chapter V.B] is a precursor to our own, but Example 4.15 shows it is not canonical. Shelah also investigated an extension of AxFri_3 axiomatizing primeness (the “primal framework”) but it is outside the scope of this paper.

We will however briefly discuss the canonicity of good frames. Good frames were first defined in [She09a, Chapter II]. We will assume the reader is familiar with their definition and basic properties. As already noted, the main difference with our framework is that a good frame is local: For a fixed $\lambda \geq \text{LS}(K)$, a good λ -frame assumes the existence of a nice independence relation \perp where only $a \underset{M}{\perp}^{\widehat{N}} N$ is defined, for a an element of \widehat{N} and $M \prec N \prec \widehat{N}$ models of size λ .

In [She09a, Section II.6], Shelah shows that, assuming a technical condition (that the frame is weakly successful), one can extend it uniquely to a *non-forking frame*:

basically an independence relation \perp where $M_1 \underset{M}{\perp}^{\widehat{N}} M_2$ is defined for $M \prec M_\ell \prec \widehat{N}$ in K_λ , $\ell = 1, 2$. For the rest of this section, we fix $\lambda \geq \text{LS}(K)$ and we do *not* assume the existence of a monster model (Hypothesis 2.1). Recall however that the definition of a good frame implies K_λ has some nice properties, i.e. it has amalgamation, joint embedding, no maximal model, is stable¹² in λ , and has a superlimit model.

Fact 6.9. If \mathfrak{s} is a weakly successful good λ -frame, then it extends uniquely to a non-forking frame (i.e. using Shelah’s terminology, there is a unique non-forking frame NF that respects \mathfrak{s}).

Proof. Uniqueness is [She09a, Claim II.6.3] and existence is [She09a, Conclusion II.6.34]. \square

As Shelah observed, Example 4.15 shows that a non-forking frame by itself need not be unique: we need to know it comes from a good frame, or at least that there is a good frame around. Shelah showed:

Fact 6.10. Assume that \mathfrak{s} is a good^+ λ -frame and NF is a non-forking frame, both with underlying AEC K . Then NF respects \mathfrak{s} .

Proof. See [She09a, Claim II.6.7]. \square

Here, good^+ is a technical condition asking for slightly more than just the original axioms of a good frame.

We also have that a non-forking frame induces a good frame:

¹²Really only stable for basic types, but full stability follows (see [She09a, Claim II.4.2.1]).

Fact 6.11. Assume K_λ has a superlimit, is stable in λ , and carries a non-forking frame NF (so in particular it has amalgamation) with independence relation (defined for models in K_λ) \perp . Then the relation $a \underset{M}{\overset{\widehat{N}}{\perp}} N$ holds iff there is $\widehat{N}' \succ \widehat{N}$ and

$M \prec M' \prec \widehat{N}'$ with $a \in M'$ so that $M' \underset{M}{\overset{\widehat{N}'}{\perp}} N$ defines a type-full (i.e. the basic types are all the nonalgebraic types) good λ -frame \mathfrak{t} . If in addition NF comes from a type-full weakly successful good λ -frame \mathfrak{s} , then $\mathfrak{s} = \mathfrak{t}$.

Proof. See [She09a, Claim II.6.36]. □

Thus we obtain the following canonicity result:

Corollary 6.12. Assume that \mathfrak{s}_1 is a weakly successful good⁺ λ -frame and \mathfrak{s}_2 is a weakly successful good λ -frame in the same underlying AEC K . Assume further \mathfrak{s}_1 and \mathfrak{s}_2 are type-full (i.e. their basic types are all the nonalgebraic types). Then $\mathfrak{s}_1 = \mathfrak{s}_2$.

Proof. Using Fact 6.9, let NF_ℓ be the non-forking frame extending \mathfrak{s}_ℓ for $\ell = 1, 2$. By Fact 6.10, NF_2 respects \mathfrak{s}_1 , so $\text{NF}_1 = \text{NF}_2$. By Fact 6.11 (the existence of a good frame implies the stability and superlimit hypotheses), we must also have $\mathfrak{s}_1 = \mathfrak{s}_2$. □

The methods of this paper can show slightly more: we can get rid of the good⁺.

Theorem 6.13 (Canonicity of good frames). Let $\mathfrak{s}_1, \mathfrak{s}_2$ be weakly successful good λ -frames with underlying AEC K and the same basic types. Then $\mathfrak{s}_1 = \mathfrak{s}_2$.

Proof sketch. Using Fact 6.9, let NF_ℓ be the non-forking frame extending \mathfrak{s}_ℓ for $\ell = 1, 2$. Let $\underset{M}{\overset{(1)}{\perp}}, \underset{M}{\overset{(2)}{\perp}}$ be the independence relations (for models in K_λ) associated to NF_1, NF_2 respectively. By Fact 6.11, one can extend their domain to allow a single element on the left hand side. Thus without loss of generality we may assume \mathfrak{s}_1 and \mathfrak{s}_2 are type-full. Let $M \prec N \prec \widehat{N}$ and let $a \in \widehat{N}$. Assume $a \underset{M}{\overset{(1), \widehat{N}}{\perp}} N_0$. We

show $a \underset{M}{\overset{(2), \widehat{N}}{\perp}} N_0$. The symmetric proof will show the converse is true, and hence that $\mathfrak{s}_1 = \mathfrak{s}_2$.

First observe that stability, amalgamation, joint embedding and no maximal model in λ implies we can build a saturated (hence model-homogeneous) model \mathcal{M} of size λ^+ . Since (as we will show) the argument below only uses objects of size λ , we can take \mathcal{M} to be our monster model for this argument (i.e. we assume any set we consider comes from \mathcal{M}). Then we have $a \underset{M}{\overset{(1), \widehat{N}}{\perp}} N_0$ if and only if $a \underset{M}{\overset{(1), \mathcal{M}}{\perp}} N_0$

so below we drop \mathcal{M} and only talk about $a \underset{M}{\overset{(1)}{\downarrow}} N_0$, and similarly for $\overset{(2)}{\downarrow}$. Note that working inside \mathcal{M} is not essential (we could always make the ambient model \widehat{N} grow bigger as our proof proceeds) but simplifies the notation and lets us quote our previous proofs verbatim.

Now, we observe that our proof of Corollary 4.14 is local-enough (i.e. it can be carried out inside \mathcal{M}). We sketch the details: First build a sequence $(M_i)_{i < \omega}$ independent (in the sense of $\overset{(2)}{\downarrow}$) over M so that $M_0 = N_0$, $M_i \equiv_M N_0$. Let $N := \bigcup_{i < \omega} N'_i$, where N'_i witness the independence of the sequence. Notice that we can take $N \in K_\lambda$, by cardinality considerations. By extension, find $f : N \equiv_M N'$ so that $a \underset{M}{\overset{(1)}{\downarrow}} N'$. Let $M'_i := f[M_i]$. By the proof of Lemma 4.11 (and recalling that $\kappa_1(\downarrow) \leq \omega$ in good frames), there is $i < \omega$ such that $a \underset{M}{\overset{(2)}{\downarrow}} M'_i$ (notice that Fact 6.11 is what makes the argument go through). Finally, use the proof of Lemma 4.7 to conclude that $a \underset{M}{\overset{(2)}{\downarrow}} N_0$. \square

We do not know whether one can say more, namely:

Question 6.14. Let \mathfrak{s}_1 and \mathfrak{s}_2 be good λ -frames with the same underlying AEC and the same basic types. Is $\mathfrak{s}_1 = \mathfrak{s}_2$?

7. CONCLUSION

We have shown that an AEC with a monster model can have at most one “forking-like” notion. On the other hand, we believe the question of when such a forking-like notion exists is still poorly understood. For example, is there a natural condition implying that coheir has extension in Fact 3.16? Even the following is open:

Question 7.1. Assume K is fully $(< \kappa)$ -tame, fully $(< \kappa)$ -type short and categorical in some high-enough $\lambda > \kappa$. Does K have an independence relation with (E) , (U) and (L) ?

Using the good frames machinery, an approximation is proven in [Bon14a] using some GCH-like hypotheses. However, the global assumptions of tameness and a monster model gives us a lot more power than just the local assumptions used to obtain a good frame.

It is also open whether such an independence relation has to be coheir (i.e. even if coheir does not satisfy (E)):

Question 7.2. Assume \perp is an independence relation with (E) , (U) and (L) . Let M be sufficiently saturated. Under what conditions does $\perp_M = \overset{\text{(ch)}}{\perp}_M$?

Finally, we note that while some of our results are local and can be adapted to the good frames context (see e.g. Theorem 6.13), some are not (e.g. Theorem 5.14, Lemma 5.4.4). It would be interesting to know how much non-locality is really necessary for such results. This would help us understand how much power the globalness of our definition of independence relations really gives us.

REFERENCES

- [Adl09a] Hans Adler, *A geometric introduction to forking and thorn-forking*, Journal of Mathematical Logic **9** (2009), no. 1, 1–20.
- [Adl09b] ———, *Thorn-forking as local forking*, Journal of Mathematical Logic **9** (2009), no. 1, 21–38.
- [Bal88] John T. Baldwin, *Fundamentals of stability theory*, Perspectives in mathematical logic, Springer-Verlag, 1988.
- [BG] Will Boney and Rami Grossberg, *Forking in short and tame AECs*, Preprint. URL: <http://arxiv.org/abs/1306.6562v9>.
- [Bon] Will Boney, *Computing the number of types of infinite length*, Notre Dame Journal of Formal Logic, To appear. URL: <http://arxiv.org/abs/1309.4485v2>.
- [Bon14a] ———, *Tameness and extending frames*, Journal of Mathematical Logic **14** (2014), no. 2, 1450007.
- [Bon14b] ———, *Tameness from large cardinal axioms*, The Journal of Symbolic Logic **79** (2014), no. 4, 1092–1119.
- [BV] Will Boney and Sebastien Vasey, *Chains of saturated models in AECs*, Preprint. URL: <http://arxiv.org/abs/1503.08781v3>.
- [GV06a] Rami Grossberg and Monica VanDieren, *Categoricity from one successor cardinal in tame abstract elementary classes*, Journal of Mathematical Logic **6** (2006), no. 2, 181–201.
- [GV06b] ———, *Galois-stability for tame abstract elementary classes*, Journal of Mathematical Logic **6** (2006), no. 1, 25–49.
- [GVV] Rami Grossberg, Monica VanDieren, and Andrés Villaveces, *Uniqueness of limit models in classes with amalgamation*, Mathematical Logic Quarterly, To appear. URL: <http://arxiv.org/abs/1507.02118v1>.
- [HH84] Victor Harnik and Leo Harrington, *Fundamentals of forking*, Annals of Pure and Applied Logic **26** (1984), 245–286.
- [JS12] Adi Jarden and Alon Sitton, *Independence, dimension and continuity in non-forking frames*, The Journal of Symbolic Logic **78** (2012), no. 2, 602–632.
- [JS13] Adi Jarden and Saharon Shelah, *Non-forking frames in abstract elementary classes*, Annals of Pure and Applied Logic **164** (2013), 135–191.
- [Kol05] Alexei Kolesnikov, *Dependence relations in non-elementary classes*, Logic and its applications (Andreas Blass and Yi Zhang, eds.), Contemporary Mathematics, American Mathematical Society, 2005, pp. 203–230.
- [KP97] Byunghan Kim and Anand Pillay, *Simple theories*, Annals of Pure and Applied Logic **88** (1997), 149–164.

- [Las76] Daniel Lascar, *Ranks and definability in superstable theories*, Israel Journal of Mathematics **23** (1976), no. 1, 53–87.
- [MS90] Michael Makkai and Saharon Shelah, *Categoricity of theories in $\mathbb{L}_{\kappa,\omega}$, with κ a compact cardinal*, Annals of Pure and Applied Logic **47** (1990), 41–97.
- [She75] Saharon Shelah, *Categoricity in \aleph_1 of sentences in $L_{\omega_1,\omega}(Q)$* , Israel Journal of Mathematics **20** (1975), no. 2, 127–148.
- [She78] ———, *Classification theory and the number of non-isomorphic models*, Studies in logic and the foundations of mathematics, vol. 92, North-Holland, 1978.
- [She87] ———, *Universal classes*, Classification theory (Chicago, IL, 1985) (John T. Baldwin, ed.), Lecture Notes in Mathematics, vol. 1292, Springer-Verlag, 1987, pp. 264–418.
- [She90] ———, *Classification theory and the number of non-isomorphic models*, 2nd ed., Studies in logic and the foundations of mathematics, vol. 92, North-Holland, 1990.
- [She99] ———, *Categoricity for abstract classes with amalgamation*, Annals of Pure and Applied Logic **98** (1999), no. 1, 261–294.
- [She01] ———, *Categoricity of an abstract elementary class in two successive cardinals*, Israel Journal of Mathematics **126** (2001), 29–128.
- [She09a] ———, *Classification theory for abstract elementary classes*, Studies in Logic: Mathematical logic and foundations, vol. 18, College Publications, 2009.
- [She09b] ———, *Classification theory for abstract elementary classes 2*, Studies in Logic: Mathematical logic and foundations, vol. 20, College Publications, 2009.
- [SV99] Saharon Shelah and Andrés Villaveces, *Toward categoricity for classes with no maximal models*, Annals of Pure and Applied Logic **97** (1999), 1–25.
- [Van06] Monica VanDieren, *Categoricity in abstract elementary classes with no maximal models*, Annals of Pure and Applied Logic **141** (2006), 108–147.
- [Van13] ———, *Erratum to "Categoricity in abstract elementary classes with no maximal models" [Ann. Pure Appl. Logic 141 (2006) 108–147]*, Annals of Pure and Applied Logic **164** (2013), no. 2, 131–133.
- [Vasa] Sebastien Vasey, *Forking and superstability in tame AECs*, The Journal of Symbolic Logic, To appear. URL: <http://arxiv.org/abs/1405.7443v3>.
- [Vasb] ———, *Independence in abstract elementary classes*, Preprint. URL: <http://arxiv.org/abs/1503.01366v5>.
- [Vasc] ———, *Infinitary stability theory*, Preprint. URL: <http://arxiv.org/abs/1412.3313v6>.
- [VV] Monica VanDieren and Sebastien Vasey, *Symmetry in abstract elementary classes with amalgamation*, Preprint. URL: <http://arxiv.org/abs/1508.03252v3>.

E-mail address: wboney@math.harvard.edu

URL: <http://math.harvard.edu/~wboney/>

MATHEMATICS DEPARTMENT, HARVARD UNIVERSITY, CAMBRIDGE, MA, USA

E-mail address: rami@cmu.edu

DEPARTMENT OF MATHEMATICAL SCIENCES, CARNEGIE MELLON UNIVERSITY, PITTSBURGH, PA, USA

E-mail address: akolesnikov@towson.edu

DEPARTMENT OF MATHEMATICS, TOWSON UNIVERSITY, MD, USA

E-mail address: `sebv@cmu.edu`

URL: `http://math.cmu.edu/~svasey/`

DEPARTMENT OF MATHEMATICAL SCIENCES, CARNEGIE MELLON UNIVERSITY, PITTSBURGH,
PA, USA